

Title: **The use and underuse of model systems in infectious disease ecology & evolutionary biology.**

Authors: Nina Wale^{1,2} & Meghan A. Duffy¹

¹ Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, Michigan USA 48109

² Department of Microbiology & Molecular Genetics and Department of Integrative Biology, Michigan State University, East Lansing, Michigan, USA 48824

Keywords: **model system, disease ecology and evolution, EEID, multiscale, within host, ecosystem**

This manuscript is submitted as part of the Vice Presidential Symposium 2020.

Data and Code Accessibility

A list of the references reviewed as part of this paper will be made available alongside the paper at datadryad.org

Abstract

Ever since biologists began studying the ecology and evolution of infectious diseases (EEID), laboratory-based ‘model systems’ have been important for developing and testing theory. Yet what EEID researchers mean by ‘model systems’ and what they want from them is unclear. This uncertainty hinders our ability to maximally exploit these systems, identify knowledge gaps, and establish effective new model systems. Here, we borrow a definition of model systems from the biomolecular sciences to assess how EEID researchers are (and are not) using ten key model systems. According to this definition, model systems in EEID are not being used to their fullest and, in fact, cannot even be considered to be model systems. Research using these systems consistently addresses only two of the three fundamental processes that underlie disease dynamics—transmission and disease, but not recovery. Further, studies tend to focus on only a few scales of biological organization that matter for disease ecology and evolution. Moreover, the field lacks an infrastructure to perform comparative analyses. We aim to begin a discussion of what we want from model systems, which would further progress toward a thorough, holistic understanding of EEID.

Introduction

Many researchers studying the ecology and evolution of infectious diseases (EEID) work on 'model systems', ourselves included. Yet, what we mean as a scientific community by a 'model system' is far from clear. The phrase 'model system' is invoked variously to mean that a particular experimental system is tractable (i.e., the host and/or parasite has a fast generation time, are easily culturable, and/or are easily manipulated (Antonovics et al. 2002; Ebert 2008; Koch and Schmid-Hempel 2011)); a model of a specific disease (Buckling et al. 1997) or studied with the goal of making some wider insight into general principles of host-parasite interactions (Antonovics et al. 2002; Huijben et al. 2018). Here, we take stock of what we do and, perhaps more importantly, do not do with 'model systems' in EEID. We do this not only to characterize the current state of model systems research in EEID, but to stimulate discussion about the conception and utility of 'model systems' in the study of infectious disease ecology and evolution.

The use, definition, and design of model systems is a matter for debate in several of the biological sciences but has perhaps received most attention in the biomolecular fields, such as genetics (Jenner and Wills 2007; Ankeny and Leonelli 2011; Katz 2016; Dietrich et al. 2019). In the biomolecular conception, model systems are a special subset of experimental systems that, first, are representative of species other than themselves and thus can be used to illuminate generalities about the focal phenomena of interest to a field (Ankeny and Leonelli 2011; Leonelli and Ankeny 2013). Second, model systems facilitate the study of multiple biological processes, from genetics to development to ecology, so that a holistic understanding of the organism is achieved (though, in actuality, the ecology and evolution of model organisms used in cellular-molecular-developmental biology is rarely studied (Alfred and Baldwin 2015)). Lastly, model systems research is ultimately motivated by the aim of performing comparative analyses in order to reveal general rules about biology (Jenner and Wills 2007; Ankeny and Leonelli 2011).

As a result, an infrastructure for the sharing of data and resources is a key component of model systems research (Ankeny and Leonelli 2011; Leonelli and Ankeny 2013). It is notable that these notions of what makes for good (model systems) research from a cellular-molecular-developmental perspective echo the statement of Joseph Travis, in his Presidential Address to the American Society of Naturalists, that ‘robust inference requires...horizontal comparisons, the examination of the same questions at the same level in a variety of systems...and vertical integration, the study of many ecological and evolutionary processes as they unfold in a single species or system’ (Travis 2006, pg. 307).

Here, we use these criteria to frame an analysis and discussion of how we use (or underuse) systems that are employed as models in EEID. We do not claim these criteria to be the ultimate criteria that define a model system but rather use them as a benchmark against which to measure our use of so-called model systems. First, with reference to these criteria, we examine how a variety of systems considered to be ‘models’ in EEID are used by researchers. Second, where we are not meeting these criteria, we discuss whether this is or is not a loss—i.e., whether these gaps actually represent ‘underuse’ of systems. Lastly, we outline points that might be discussed so that the field can shed the definition of model system that scaffolds our review and build a bespoke definition for EEID.

Part I: The use of 'model systems' in EEID

To establish the extent to which the experimental systems in EEID are models *sensu strictu*, we conducted a review of the literature involving 10 experimental systems (Table 1).

Methodology

Key Phenomena

According to the definition of a model system outlined above, one distinguishing feature of a model system is that it is an experimental system in which the fundamental processes of interest to a (sub)discipline can be and are examined. So, to understand the extent to which experimental systems used by disease ecologists and evolutionary biologists are 'model systems', we need to first establish what processes or phenomena are of fundamental interest in EEID.

Theory provides an answer, by formally delineating the processes that drive disease dynamics. The vast majority of theoretical studies of infectious disease dynamics begin with some form of Anderson and May's models of disease dynamics (Anderson and May 1979; May and Anderson 1979). According to these models, a healthy, susceptible (S) host becomes an exposed (E) and then infectious (I) host by the process of pathogen transmission. During either of the infected states (exposed or infectious), hosts may experience disease. Hosts then exit the infected states via the process of recovery (or, in some cases, death) (Fig. 1A).

Both pathogen (Fig. 1A pink) and host traits (Fig. 1A, blue) influence transmission, disease, and recovery. In the case of transmission, whether a host becomes infected is dependent on, on the one hand, the pathogen's infectivity and, on the other, the host's susceptibility to infection. Once inside the host, parasites may replicate, so that the parasite population grows and/or develops to a new stage; for simplicity we term both of these 'replication' in our review. In the final stage of the infection, parasites may produce transmission stages (e.g., eggs or gametocytes, in the case of helminths and malaria parasites, respectively), which disperse from the primary host to a secondary host. The amount of disease a host experiences when it is infected depends on the virulence of the pathogen, here defined as the parasite's contribution to disease, and immunopathology, when the host's immune response itself causes harm; the host's ability to minimize the damage associated with a given parasite burden is termed tolerance (Little et al.

2010). Lastly, recovery occurs either because the infection is self-limiting or because of the host's resistance, defined here as the capacity of the host to kill or expel pathogens (rather than as the opposite of susceptibility, as it is sometimes defined e.g. Thrall and Burdon 2003).

The parasite and host traits that govern transmission, disease, and recovery can vary between individuals, be heritable and have significant fitness effects; thus, these traits can and do evolve in response to selection. Both parasite and host traits are subject to selection from ecological factors such as abiotic or biotic stressors (e.g. predators); moreover, host phenotype exerts selection on parasite traits, and vice versa. Indeed, we outline the traits involved in each phenomenon in sequence and as pairs of host and parasite traits to emphasize the fact that coevolution occurs and can happen at several stages of the infection process (Duneau et al. 2011). In addition, at the level of the host or parasite population, the presence and strength of trade-offs between the traits that govern the different phenomena influence the trajectory of trait evolution. For example, the virulence-transmission trade-off that lies at the heart of models of parasite evolution (reviewed in Cressler et al. 2016) is determined by the relationship between within-host replication, virulence, and dispersal capacity. Similarly, trade-offs between host resistance and tolerance can influence the evolution of these traits (Raberg et al. 2007) and, ultimately, the amount disease associated with a particular infection.

In our literature review, we noted which of the traits that contribute to the key phenomena in EEID (Fig. 1A) were being studied. In some studies, we were unable to isolate which of the transmission traits were being studied and/or multiple traits were assessed. For example, in a study of epidemic dynamics, the focus is not on parasite replication *or* infectivity *or* dispersal capacity but rather on the manifestation of these processes (what we call 'realized transmission'). In such cases, we used other terms, the definition of which can be found in Table 2, to classify what aspect of transmission was being studied. Similarly, many studies were not

designed to tease apart the impact of (heritable variation in) virulence and tolerance on disease — i.e., they did not examine how disease changed when parasite strain was varied in a single host genotype or when hosts of different genetic backgrounds were exposed a single parasite strain. We thus recorded only which manifestation(s) of disease (morbidity, mortality, etc.) were being monitored. Lastly, EEID researchers sometimes study other impacts of infectious disease, for example parasite nutrient content or the positive effects of parasites on hosts. These studies were allotted to an ‘other’ category. As we conducted our review, we tracked which of the traits that contribute to transmission, disease, and recovery were studied, how they were measured, and were alert to trends in research methodologies.

Focal questions in EEID & the scales of biological organization at which they are studied

Disease ecologists and evolutionary biologists are interested in the factors that drive variation in transmission/disease/recovery among individuals and populations, as well as the consequences of this variation (Box 1). The ecological interactions that impact trait variation occur at a variety of scales of biological organization (Lively et al. 2014); in turn, infectious diseases can impact ecological interactions at a variety of scales (Fig. 1b). Therefore, in addition to noting which of the key phenomena each study addressed, we also recorded the scale of the phenomenon/trait being measured (i.e., the scale of the dependent variable) and the scale at which the variation was generated (i.e., the scale of the independent variable in the experiment). For example, in a study of the impact of population demography on the susceptibility of hosts, we recorded that the dependent variable (host susceptibility) was at the individual level and the independent variable (population demography) was at the host population scale. Table 3 and Fig. 1b contain details of which experimental variables were assigned to which scale.

Systems reviewed

We examined which of the key phenomena (i.e. transmission, disease, recovery) were addressed using a variety of experimental systems (Table 1). In selecting these systems, we attempted to cover the history of the field, and a diversity of parasite types, from viruses to helminths, and host taxa, from plants, to invertebrates, to mammals. So as to cover as great a breadth as possible, we set two criteria determining whether a system could be included. The first was that researchers had to have the ability to induce infection experimentally, since the capacity to manipulate the presence of infections seems to be a basal requirement for their experimental study. We did not require that the entire transmission cycle of the parasite could readily be completed in the laboratory since, as we discuss more in the “Population & Epidemic Level” section below, doing so would have severely restricted our ability to conduct this review. Second, the system had to be a focus of ecological/evolutionary study for greater than ten years, since researchers cannot be expected to have established a holistic study system in less time than that. Beyond that, we endeavored to limit overlap between host or pathogen types.

By necessity, the list of systems (Table 1), though broad, is not exhaustive. First, we omitted a number of important pathosystems where experimental inoculation of the host is possible but is used relatively rarely as an experimental methodology (at least among the studies recovered using our search criteria). This was the case for the snail-trematode system *Microphallus-Potamopyrgus*, for example, which has made an important contribution for our understanding of host-parasite coevolution (Dybdahl and Lively 1998; Lively and Dybdahl 2000; Morran et al. 2011). Second, we omitted bacteriophage-bacteria systems, though they permit experimental inoculation of hosts and have made a substantial contribution to our understanding of host and parasite (co)evolution and pathogen emergence (Horne 1970; Chao et al. 1977; Lenski and Levin 1985; Benmayor et al. 2009; Koskella 2014). Given the myriad host strain-phage combinations utilized in this subfield, this literature was unwieldy to review; it deserves a standalone review that can also address the specific biology of virus-bacteria interactions. In

addition to systems where experimental inoculation is possible, there are several pathosystems where infections are not experimentally generated routinely but nonetheless have made important contributions to our understanding of disease dynamics (Hudson and Newborn 1998; Ezenwa and Jolles 2015), as have observational studies or field experiments in systems we did review, such as those involving *H. polygyrus* (Gregory et al. 1990; Knowles et al. 2013). We recognize that, even within the inclusion criteria established above, there are additional systems that could have been included (e.g., the stickleback-*Schistocephalus* system (Barber and Scharsack 2010), the *Silene-Ustilago* system (Alexander 1989)) but nevertheless believe the diverse systems that are included well reflect dominant research trends in EEID.

Literature search protocol

We searched for scientific literature related to each system using Web of Science (Thomas Reuters). We searched for the scientific name of the pathogen, including nomenclature that had been previously used but since abandoned - for example, in the case of the *Heligmosomoides polygyrus*, we also searched for *Nematospiroides dubius*. We identified the 10 most cited and 5 least cited studies that included at least one laboratory-based experiment in which hosts were exposed to the pathogen (even if that was the inoculation of hosts prior to their distribution into the field or laboratory-based phenotypic assessment of field collected material). We included least cited papers not only because they were likely to include the most recent research but also because personal experience has taught us that these papers contain some of the oldest and/or most idiosyncratic research, and so could reveal ideas once pursued in EEID but since abandoned. In the case of some pathogens that are also important model systems in other fields, e.g., the rodent model of poxviruses, infectious ectromelia virus, and the rodent intestinal roundworm, *Heligmosomoides polygyrus*, the top ten/bottom five cited papers did not address ecological and evolutionary questions. When this occurred, we performed an additional search

to identify papers of an ecological or evolutionary nature, by searching for 'ecolog* OR evolution*' in the topic field in addition to the pathogen species name.

Lastly, to further capture older research, we identified one scientist who pioneered the study of each model system that we reviewed (Table 1, we list several researchers for completeness but only the research of the bolded researcher was searched for). When fewer than five papers by the 'founder' appeared in the first search, a second search was conducted with the system name(s) and the name of the scientist. We added the most cited references that appeared in this secondary search to our primary search results, leading to a total of five papers by the 'founder' in the final collection of papers for the system. In the case of infectious ectromelia virus, it was clear that Web of Science did not retrieve historical references since it uncovered some, but not all, of a set of research articles conducted by the system's 'founder' Frank Fenner, which were numbered in sequence. We thus searched for this author's bibliography directly, to recover the missing papers from this sequence. That Web of Science did not uncover these older references highlights that there are limitations associated with using this (or indeed any other) search engine for this review. As the field of EEID became distinct from its relatives around the time that Anderson and May published their landmark papers, and the majority of systems reviewed here were established as experimental systems at that time or afterward (Table 1), we do not think this greatly impacts our conclusions, however.

With the papers collated, we then extracted the following information from each paper: i) which of the umbrella phenomena (transmission, disease, recovery) was investigated, ii) the variable(s) manipulated, iii) the variable(s) measured, and the scale at which these variables were measured or manipulated (see '*Focal questions in EEID & the scales of biological organization at which they are studied*', above).

Results

We present the results of our review with a focus on the extent to which those systems used as models in EEID address the key phenomena of interest to the field — transmission, disease and recovery — and the extent to which we study them at the biological scales at which disease dynamics occur. As such we seek to understand the extent to which model systems are studied holistically and are representative of the processes EEID researchers are interested in. Where possible, we highlight specific features of the different systems that enable disease ecologists and evolutionary biologists to study certain phenomena in them.

1. Are we studying the key phenomena in EEID?

All of the key phenomena — transmission, disease, and recovery — are represented in the studies reviewed. However, the extent to which all three of the phenomena are addressed varies among systems, as does the extent to which they are used to study the traits that drive these key phenomena (Fig. 2a).

Transmission

Transmission was the most studied phenomenon, being the focus of more than 80% of the 160 studies reviewed. However, the parasite traits involved in transmission (Fig. 1A) were not equally represented (Fig. 2a). Infectivity and replication/development were well represented, being the focus of approximately 25% and 30% of the studies of transmission, respectively. However, only ~10% of the studies reviewed focused on the capacity of parasites to exit the host, for example via the production of transmission stages. A similarly small proportion followed the infection from its initiation to onward transmission (“realized transmission”). Although realized transmission was studied in 7/10 pathosystems, it was only a regular focus of study (i.e., the focus of $\geq 20\%$ of the papers) in four systems: IEV, *C. bombi*, *D. spathaceum*, and *R. ondatrae*. Notably, the system in which realized transmission was most often studied — IEV —

is no longer in use as an experimental system in EEID, somewhat ironically because it was so transmissible as to be considered a biohazard (Fenner 2000)! Rather than measure transmission directly, most researchers took pathogen load as a proxy for it.

In addition to parasite traits, transmission also involves host susceptibility, narrowly defined here as the capacity of a host to become infected. Of those reviewed, *M. lini* was the system most used to study susceptibility, which is fitting given its role in the development of the gene-for-gene model of susceptibility/infectivity (Flor 1971). The genetic component of susceptibility and how it varies in space and time was the major focus of studies employing *M. lini* and *P. ramosa*, which also coevolves with its host through the matching of genes that mediate infectivity/susceptibility (Thrall et al. 2016). Interestingly, our search did not uncover studies of recovery/resistance in *M. lini* and *P. ramosa*. This could imply that there is either an evolutionary trade-off between defenses that block and clear infection (why invest in resistance if you don't get infected, after all?) or, recognizing that we reviewed only a subset of papers published in each system, this might reflect the relative degree to which different topics have been studied and/or cited. Indeed, we know there are studies in *P. ramosa* that have focused on post-infection resistance (e.g., Hall and Ebert 2012). In addition to the genetic basis of susceptibility, the behavioral component of host susceptibility was also examined, primarily in the *D. spathaceum* and *C. bombi* systems (Hernandez and Sukhdeo 1995; Karvonen et al. 2004; Gegear et al. 2006; Bouwman and Hawley 2010; Milotic et al. 2017; Fouks et al. 2018; Mikheev et al. 2019).

Disease

The negative impact of infection on host fitness, which we define as disease, was the focus of more than half of the studies that we assessed and was investigated in all systems.

Surprisingly, despite their conceptual importance in disease ecology, fewer than half of the

studies of disease focused on virulence and tolerance. Disease is a manifestation of the interaction of the host and parasite: here, we use “virulence” to denote the parasite’s contribution to disease, and “tolerance” to denote the amelioration of that effect by the host. Approximately a sixth of the studies of disease focused on virulence, fewer than 5% on tolerance. Studies of virulence were conducted in five systems. Of these, genetic variation at the level of both the host and the parasite was exploitable for laboratory studies in two systems, *M. gallisepticum* and *P. chabaudi*. This permitted the quantification of heritable variation in tolerance and virulence to disease in the same system, a necessary precursor for studying the coevolution of these traits (though it is questionable how relevant that is in the *P. chabaudi* system, given that the host and parasite do not naturally cooccur) (Little et al. 2010). The majority of studies of disease did not focus on the contributions of host and parasite genetic variation to variation in disease severity. Rather, these studies focused on describing the symptoms of infection (e.g., Weimer et al. 1955; Goodman and Johnson 2011) or how they change with the host environment (Ferguson and Read 2002; Paull and Johnson 2011; Overholt et al. 2012).

Disease was measured using qualitative and quantitative measures of varying ecological and evolutionary significance. In some systems, like *P. ramosa* — which is a castrating obligate-killer of short-lived and rapidly reproducing zooplankton — the impact of infection on survival could be directly assessed (Ebert 2008). In others, disease was measured via quantitative health metrics such as cellular numbers, as in *P. chabaudi* (Taylor et al. 1998), or via a qualitative metric, as in *M. gallisepticum*, where the severity of conjunctivitis in birds was measured via an ‘eye score’ (Kollias et al. 2004). In relatively long lived animals, such as mice, fish, and birds, it is difficult to know how these measures translate into life-long fitness (Graham et al. 2010), though our search did uncover studies where symptom severity was related to the likelihood that hosts were predated upon (Seppälä et al. 2004, 2005). Furthermore, the effect of

infection on behaviors directly related to fitness, such as mating (Kavaliers et al. 2003), foraging (Crowden and Broom 1980; Ponton et al. 2011; Adelman et al. 2015) and predation (Webster et al. 1999), was assessed in a number of systems. Notably, in the studies we reviewed, the effect of infection on hosts' susceptibility to predation was only assessed in systems where parasites are trophically transmitted (*D. spathaceum* and *H. diminuta*) and, as such, these studies were as much studies of the fitness of the parasite as the host.

Recovery

Recovery was the least well studied of the three key phenomena, the focus of just 16% of the studies, although it was studied in the majority (7/10) of systems. In part, the paucity of studies in this category is likely to be a consequence of the fact that we assigned only a single trait, resistance, to it. We define resistance as the capacity for hosts to kill or eject parasites. Notably, hosts routinely clear parasites (and hence are able to 'recover completely') in only 4/10 of the studies systems reviewed: IEV, *H. polygyrus*, *P. chabaudi*, and *M. gallisepticum* (Table 1).

In the vast majority of studies, resistance was measured by quantifying the reduction in parasite load after the infection was established. As such, whether the host was involved in the clearance of parasites (per the above definition of resistance) or whether parasite populations were self-limited (which itself could be host-induced (Hite et al. 2019; Wale et al. 2019)) was not always delineated. The host immune response to infection was measured directly in *H. diminuta*, *H. polygyrus*, *D. spathaceum*, IEV, *M. gallisepticum*, and *P. chabaudi*, the systems in which the host response can be manipulated (usually by pre-exposing hosts to the pathogen). It is likely that such 'mechanistic' studies of resistance/recovery are possible in these systems because they involve vertebrate hosts whose immune system is relatively well characterized. Interestingly, though one of the oldest studies of *D. spathaceum* is mechanistic in nature (Stables and Chappell 1986a), and the system was established by parasitologists/immunologists, it was not used for

mechanistic studies of the host response in any other study reviewed. Rather, *D. spathaceum*, along with *C. bombi*, was used to investigate how factors beyond the individual host impact the host's capacity to control infections, including abiotic factors (Stables and Chappell 1986b; Palmer-Young et al. 2019), social group-composition (Klemme and Karvonen 2018) and socially transmitted microbes (Koch and Schmid-Hempel 2011; Koch and Schmid-Hempel 2012); however, these studies were not included in the recovery category, because they do not address inherent host resistance.

We shaped our review around an SIR framework, which was formulated to describe the dynamics of parasite transmission, and hence our discussion of recovery focuses on the extent to which hosts bear parasites (and so can transmit them). However, we recognize that other processes might be reasonably included in a discussion of recovery e.g. tolerance or post-infection disease tissue repair. That said, even had we included these processes in the category of recovery we might still have found that it receives little attention. For studies of tolerance were rare among those reviewed (see above) and we recall no studies in which hosts were followed up post-infection.

At what scales are we studying key phenomena?

Approximately a third of the studies involve researchers looking at how 'individual level' factors — e.g., parasite strain (Fenner 1949a; Rodriguez and Kleven 1980), host age (Fenner 1947a) or immune status, size of parasite inoculum (Fenner 1947b; Johnson et al. 2001; Karvonen et al. 2003), or inoculation method (Fenner 1947b; Lepak and Thatcher 1962) — impact individual level variables. Since the majority of studies do not focus on interactions between individual-level processes, we focus on other scales from here on. We first address each scale in its capacity as a dependent and then an independent variable; some studies appear in multiple

categories. To stimulate future research, we identify challenges involved in doing research at these different scales.

Within-host

Within-host processes were either manipulated or measured in just over a third of the studies reviewed. The dynamics of parasite populations and their distribution within the host were the subject of the earliest papers that our review uncovered regarding *D. spathaceum* (Betterton 1974), IEV (Fenner 1947a), *H. diminuta* (Chandler 1939), and *H. polygyrus* (Dobson and Owen 1978). Within-host dynamics were the response variable in approximately a quarter of all studies reviewed. These addressed the way that within-host dynamics changed with parasite population parameters, such as initial size and genetic background, and host factors such as immunity, diet (Bansemir and Sukhdeo 1996; Ponton et al. 2011), and even temperature (Stables and Chappell 1986a).

A major theme of studies in the within-host processes category was interactions among parasites. The majority of experiments we reviewed focused on intraspecific interactions and interrogated the (dis)advantage of focal parasite traits, including virulence, drug resistance, and motility, in different host environments. Variability in host environments (and hence within-host interactions) was created by modulating intrinsic host traits such as host sex (Gipson et al. 2019) or via the administration of medications (Wargo et al. 2007; Huijben et al. 2015). Interspecific interactions were similarly, though less often, explored (Holmes 1959; Graham et al. 2005; Lass et al. 2013; Budischak et al. 2015). Some of the more recent studies in the corpus of papers we reviewed focus on the interaction between parasites and the microbiota, rather than on parasite-parasite interactions (Koch and Schmid-Hempel 2011; Koch and Schmid-Hempel 2012; Clerc et al. 2015; Zaiss et al. 2015). Interestingly, these papers were often focused on the use of parasites to alter microbiota to promote host fitness; in contrast, of

the papers in our review that focused on parasite-parasite interactions, the predominant focus was on parasite fitness.

Within-host processes are often dynamical in nature and thus necessitate the tracking of host or parasite populations within the host's body. In the majority of study systems, the dynamics of within host populations were tracked by the sequential culling of cohorts of animals. The exceptions were *M. gallisepticum* and *P. chabaudi*, in which non-destructive, repeated sampling of parasite and disease parameters (conjunctivitis and anemia, respectively) from single individuals was possible. This sampling methodology is made possible by the spatial distribution within the host of these pathogens, which fall at extreme ends on a gradient of tropism:

Plasmodium chabaudi is distributed in the blood (at least during the asexual stage of infection), so one can sample the periphery without destroying the animal; *Mycoplasma gallisepticum*, by contrast, is concentrated in and around the eye (though it can be found elsewhere (Dhondt et al. 2005)).

Population & Epidemic Level

Laboratory studies rarely focus on epidemiological dynamics, either as an independent or dependent variable; only 14% of the studies reviewed address disease dynamics at this scale. This implies that the rarity of studies of realized transmission at the individual level might be due to difficulties with the initiation of transmission in the laboratory, rather than to the relative ease of measuring parasite load as compared to realized transmission. Indeed, in half of the systems the entire transmission cycle of the parasite cannot be readily completed in the laboratory (Table 1), ruling out the study of epidemic dynamics and their subsequent impacts on parasite and host population dynamics.

Two systems dominate the study of epidemiological dynamics, IEV and *C. bombi*. IEV is the focus of the earliest epidemiological investigations in our review, which focus on describing how ‘individual’ level variation of host and parasites, such as strain, age, and immunity, impact epidemics (Fenner 1948a, 1948b, 1949b). By contrast, those studies involving *C. bombi* focus on the impact of features of the host population, specifically density (Bailes et al. 2020), structure (Otterstatter and Thomson 2007), relatedness (Shykoff and Schmid-Hempel 1991), and turnover (Buechel and Schmid-Hempel 2016), on epidemic dynamics. Experiments that utilize *C. bombi* involve many more replicates than those using IEV, presumably because they involve smaller, invertebrate organisms whose populations can be established in replicate without as much maintenance or as many ethical concerns. This feature also makes epidemic level studies possible in the *P. ramosa* system (Ebert et al. 2000). Notably, the evolutionary impact of epidemics was not examined in any of the studies we reviewed.

Importantly, in several of the systems reviewed, laboratory studies were conducted alongside studies of population/epidemic scale processes in the field. Such field studies facilitate the study of (co)evolutionary dynamics (Thrall et al. 2002; Ebert 2008; Bonneaud et al. 2018) that, in turn, further laboratory investigations. For example, approximately half of the studies of the effect of population variability on individual-level infection traits were conducted using *M. lini* and *M. gallisepticum*. In each of these systems, spatial dynamics of epidemics in the field drove parasite and host evolution that was then exploited to study the heritable variation in infectivity, virulence, and immunity in the laboratory (Thrall and Burdon 2003; Hawley et al. 2013; Bonneaud et al. 2019). *P. ramosa* provides similar opportunities to sample a diversity of pathogens and hosts, as well as to analyze the effect of epidemics on host evolution (Duncan and Little 2007).

Community level

The relationship between interactions at the community level and the dynamics of infectious diseases comprised ~10% of the studies reviewed. With the growing importance of zoonotic diseases as a public and animal health problem, how parasites move between multiple hosts, and why some hosts support pathogens, but others do not, has been the focus of much recent research (Johnson et al. 2012, 2019; Mollentze et al. 2014; Fenton et al. 2015; Olival et al. 2017). It was surprising, then, that only 6% of the studies reviewed examined variability among infections in different host species. The question of why some hosts are 'better hosts' than others was specifically addressed in *H. diminuta* (Read and Voge 1954; Johnson et al. 2012) and *R. ondatrae* (Johnson et al. 2012); the former study was one of the oldest of all those reviewed. As an aside, the variation of infections among different host *genotypes*, which might be considered a topic analogous to host-species variation, was a focus of a further ~10% of studies reviewed (these were not counted as 'community' level studies). Of these, two studies distinguished themselves by using experimental evolution rather than a standing trait variation to study host range (Dobson and Owen 1977; Brindley and Dobson 1981).

In addition to studies focused on the effect of host species identity on disease dynamics, further studies investigated how the community of hosts, specifically its composition, altered disease dynamics (Johnson et al. 2013, 2014). There was only one study in which the impact of parasites on the formation or function of ecological communities (as opposed to the other way around) was the focus. Specifically, (Yan et al. 1998) investigated the impact of *H. diminuta* on interspecific competition between beetles. Other community-level studies focused on the interaction of parasites and predation (likelihood) in trophically transmitted systems (Seppälä et al. 2004; Orlofske et al. 2012).

There were a few studies that investigated how parasites withstand the ecological environment outside of their host. Of these, two studies focused on the vulnerability of the parasite to

predation (Morgan et al. 1997; Orlofske et al. 2012), while the remainder focused on abiotic factors, including light, temperature, humidity, and solar radiation (Flor 1958, 1960; Voge and Heyneman 1958; Voge 1959a, 1959b; Overholt et al. 2020; Rogalski and Duffy 2020). Those studies that were of an ecological nature — Flor’s studies of the impact of X-ray radiation on *M. lini* were not motivated by an interest in radiation *per se* but in the mutations that resulted from it — were conducted using *H. diminuta* and *P. ramosa* and were, respectively, some of the oldest (1950s) and newest (2019-2020) studies reviewed. It is likely that studies of parasites in the environment are a rarity in our review because EEID researchers are often most interested in *interactions* between host and parasite, rather than in parasites alone. Since the survival and reproduction of parasites outside of the host can be a crucial determinant of parasite fitness and transmission, however, this area deserves attention.

Ecosystem level

Given the lack of epidemiological-level studies and that we were focused on studies with at least one laboratory-based experiment, it is perhaps unsurprising that few studies address ecosystem-level impacts. Just two of the studies reviewed, both of which involve *R. ondatrae*, can be considered to be truly focused on ecosystem processes: one examines the impact of elemental nutrient supply on pathogen spread (Johnson et al. 2007), the other the nutritional value of individual parasites for consumers (McKee et al. 2020). Our review also uncovered another study of the chemical/nutritional analysis of parasites, conducted over four decades before the latter and used the *H. diminuta* system (Roberts 1961). However, the results of this study — though eminently relevant to ecosystem processes — were interpreted in the context of their relevance to host-parasite interactions within-host rather than to ecosystem processes.

To what extent are we studying our systems holistically?

A goal of model systems research *sensu strictu* is to study all of the key phenomena that drive the focal process of interest. Disease dynamics are not only influenced by ecological interactions that occur at multiple scales (within-host, epidemic, etc.; Fig. 1B) but interactions that occur at one scale may have a dramatic impact on those at another. However, our review suggests that the effect that processes at different scales have on one another is rarely studied.

There are two important trends regarding this lack of focus on interactions of processes at different levels. First, there is an absence of studies on how within-host or individual-level variation impacts population and ecosystem level processes (Fig. 3, bottom left of panels). Similarly, studies rarely examine how ecosystem- or community-level variation impacts population- or ecosystem-level processes, respectively. These trends are unsurprising given the rarity of studies at the population/epidemic level, as discussed above, and the overall lack of work on infectious diseases at the ecosystem-level (Preston et al. 2016).

Though it is uncommon to study how processes at ‘small’ scales impact ‘large’ scale processes (and vice versa), processes at multiple scales are varied in most systems (i.e., the top two rows of fig. 3 are often filled in many cells left to right, at least up to ‘population’ processes). It is most common for processes within hosts, at the individual level, and the host environment to be varied within a single system. Presumably, this is because these scales can be varied using individual hosts in the laboratory (e.g., one could inoculate an individual host with multiple parasites or remove parasites using antimicrobial drugs, vary the strain of parasite, and alter the husbandry of hosts). The consistent focus, across systems, on the (interactions between) small scale processes suggests that there is significant potential for horizontal integration *sensu Travis* in this area.

So, to what extent are our experimental systems model systems *sensu strictu*?

Our review of the most and least cited studies of the dominant model systems in EEID revealed that these ‘model systems’ are mere ‘experimental systems’ — i.e., they are used to investigate particular phenomena or processes as opposed to systems used to establish a holistic understanding of the EEID (Ankeny and Leonelli 2011). While we address both transmission and disease in the laboratory, the third key process of interest to EEID, recovery, is often left unaddressed, not least because we often use systems in which it does not occur. In terms of the scale of interactions that these experimental systems are used to explore, researchers succeed in building a holistic understanding of host-parasite interactions at the level of the individual host (or parasite strain). Ironically for a field founded in population biology, however, we fail to scale up these studies to that of populations and above, at least in the laboratory.

Part II: The underuse of ‘model systems’?

In the previous section we used the definition of model systems borrowed from the biomolecular sciences to investigate how we use experimental systems in EEID and, by extension, how they are not used. To understand whether the gaps identified should be filled, we must assess whether these gaps hold back our understanding of EEID and, if so, if and how they could be remedied. That is, we must assess the extent to which our experimental systems are indeed underused.

The transmission gap

Since onward transmission is the ultimate realization of pathogen fitness, its relative rarity as an experimental outcome is notable. However, at least in the case of individual level studies their relatively rarity may not amount to an omission. Experimental proxies of onward transmission can be an appropriate measure of parasite fitness and allow for experiments with large numbers of individuals. In order for proxies to be of maximal use in the analysis of parasite fitness, these proxies must be measured at the level of the individual parasite-host strain combination (e.g.

Bell et al. 2012) and in a context that reflects natural transmission, for measures of parasite fitness such as parasite growth rate and virulence change greatly in a non-natural context (Fenner 1947*b*; Spence et al. 2013).

The major limitation of using systems where the parasite cannot complete its life cycle in the laboratory (Table 1) is that it reduces the capacity to study epidemiology experimentally, something that was once a priority of EEID and arguably deserves concerted investment. Experimental studies of the epidemiology of IEV were important in the conception of the Anderson and May's classic models of infectious disease dynamics (Anderson and May 1979). Around the same time that those classic models were published, Anderson and colleagues led a push to develop new systems with which to study experimental epidemiology, including the *H. diminuta* (Keymer and Anderson 1979), *Gyrodactylus* sp. (Scott and Anderson 1984), and *H. polygyrus* (Keymer 1985) systems. Indeed, in 1985, Keymer wrote a manifesto of sorts, outlining why epidemiological study systems were required and what they should look like (Keymer 1985) — arguments that stand the test of time and resonate with the arguments made herein. She argued that experimental epidemiological systems must allow the collection of data on 'the complete epidemiological behavior of the parasite in its host population' (i.e., epidemic level data), and 'the experimental study of population parameters ...their dependence on, and interactions with other biological processes and...physical variables' (pg. 56) (i.e., individual, ex-host level data). That is to say, with such a system, one can isolate the effect of factors predicted to influence epidemic dynamics (e.g., host genetic variability), tease apart how combinations of factors manifest as epidemiological dynamics, and model variation that does not (yet) appear in nature. Indeed, experimental epidemiological studies could represent an additional, complementary source of high-quality data that could be used to widen the canon of theoretical epidemiology, which is dominated by diseases that are viral, acute, and often target juveniles (Keeling and Rohani 2011). As such, experimental epidemiology can not only help to

reveal the mechanisms underlying the dynamics we observe in nature but illuminate what could be possible in nature, in all its diversity.

Filling the transmission gap could boost 'vertical integration' in EEID

A further advantage of boosting the capacity to study epidemics in the laboratory is that it will permit the mechanistic investigation of feedbacks between 'small scale' within-host processes and 'larger-scale' epidemiological processes; that is, it will fill the bottom left of the heatmaps in Fig. 3. The study of interchange between processes at these scales has been the subject of much recent theoretical investigation but is difficult to achieve in the field (Day et al. 2011; Mideo et al. 2013a, 2013b). Take the evolution of drug resistance as an example. Recent theoretical studies have suggested that, depending on the prevalence of circulating strains, within-host competition might alter the selective benefit of using different drug-dosing regimens (Hansen and Day 2014). Manipulating dosing regimens and/or epidemiological parameters can be unethical and extremely difficult in human populations, where there are also considerable challenges distinguishing between *de novo* or acquired resistance. An experimental system in which one could track the population dynamics of parasites within-host, as well of those of hosts and parasites at the epidemic scale, would be invaluable in testing and expanding this theory.

The development of systems in which we can study the interactions and feedback between processes at the within-host and population scale (and even at larger scales) may be inherently difficult, however, because the traits that make it good for one come at the disadvantage of the other. On the one hand, to facilitate the study of epidemic dynamics, hosts must be small enough that replicate, dense populations can be established, housed long-term, and manipulated. On the other, the study of within-host dynamics over time in individual hosts requires that hosts be robust enough to be repeatedly handled and sampled. Pathosystems of large insects of agricultural importance may fall into this crucial 'sweet spot'. Indeed, early work

with *Tribolium* contributed to the understanding of the dynamics of host and parasite populations (Park 1948; Keymer and Anderson 1979), and they have recently been established as an ecological-evolutionary-immunological model (Tate and Graham 2017; Jent et al. 2019). As Keymer pointed out years ago (Keymer 1985), vertebrate models of human disease are well-placed for this work, since much is known about vertebrate immunity; the development of tools to non-invasively track infection dynamics (e.g. quantitative PCR, *in vivo* imaging) and manipulate parasite and host genetics since then, have only added to their advantages. (Keymer 1985). Recent studies in which laboratory mice were 'rewilded' in order to understand the relative contribution of environment and genetics to within-host interactions (Lin et al. 2020; Yeung et al. 2020) demonstrate the unique utility of laboratory mice for answering questions in disease ecology. Such studies might be extended to study the interchange between within-host and epidemic dynamics, if technologies that permit cheap, regular, high-throughput sampling of individuals could be developed. These studies also demonstrate that leading model systems in the wider biological sciences might also be powerful systems for EEID research.

While it is clear that laboratory studies could be better harnessed to understand disease dynamics at or below the epidemic scale, it is not immediately obvious that they could illuminate epidemic processes at the scale 'above' epidemics, especially the ecosystem level. Such experiments would necessitate the use of micro- or meso-cosms e.g. per Johnson et al. (2007). However, the suitability of these types of experiments for the study of ecosystem-level processes has drawn repeated criticism because often even very large mesocosms are incapable of capturing the spatial or temporal scale of the processes of interest, the diversity of organisms involved (particularly those at higher trophic levels), or physical/chemical structure of environments (Carpenter 1996; Schindler 1998). It may thus be best to reserve mesocosm-scale experiments for estimating parameters of mathematical models (e.g. the biomass of pathogens or the (change in) nutrient content of infected hosts) or to use them only where

ecologically-relevant effects are likely to be observed e.g. in systems that involve pathogens of ecologically dominant hosts species and simple, linear food webs (Mitchell 2003; Duffy 2007; Borer et al. 2009).

Toward the cryptic & rare

Our review suggests that we are not using our model systems to their full potential for the study of interactions between parasites and host immunity and their contribution to eco-evo dynamics of disease. Of the studies we reviewed that addressed infections that do not resolve, few used measures of infection ‘success’ that permit the investigation of host defenses other than those that make the host refractory to infection. For example, in studies of *D. spathaceum*, *P. ramosa*, and *R. ondatrae*, parasite ‘infectivity’/host ‘susceptibility’ was often measured as the proportion of hosts who bore parasites at the site from which the parasites leave the host. This measure confounds the process of parasite infectivity with the processes of development and migration and, on the other hand, suggests that there is no host defense other than that which prevents infection in the first place. However, host responses that destroy parasites after the parasites successfully penetrates the host are present in these systems (Hall and Ebert 2012; LaFonte and Johnson 2013). Methods for explicitly measuring a parasite’s progress through the host e.g. imaging of fluorescent parasites (e.g., Duneau et al. 2011; LaFonte and Johnson 2013) and mathematical models that can quantitate how the hosts resist and tolerate infection (e.g. (Wale et al. 2019) could refine our understanding of which traits mediate parasite and host fitness at each stage of the infection (Duneau et al. 2011). Indeed, there is much to gain from the methodologies used in cellular-molecular biology, such as fluorescence-based microscopy and cytometry, as well as that field’s mechanistic understanding of host-pathogen interactions. By harnessing these methods and insights, we can increase our ability to observe, quantify and manipulate the phenotypes of interest to EEID, as well as promote the vertical integration of infection biology, as a whole.

There is work to do in systems where hosts can recover from infections too, specifically in the area of chronic infections. Of those infections where hosts can recover, low level chronic infections of multiple weeks to years can develop in at least some animals (Hawley et al. 2005; Achtman et al. 2007; Sakala et al. 2015). Yet we recovered few studies in which these chronic infection dynamics were the subject of study, despite the fact that chronic infections can make substantial contributions to the maintenance of disease in an epidemic context and alter the measured shape of virulence-transmission relationships. No doubt, the rarity of chronic infection studies is because, by definition, they take time. We must be careful not to take shortcuts however, as our cousins the immunologists learned, when attempts to accelerate the development of chronic infections by using larger pathogen inoculum than was normal led to a qualitatively different immune responses and so defeated the whole exercise (Vidlak and Kielian 2016).

While studying only the immunological phenomena that are easy to observe could skew our understanding of the ecology and evolution of infectious diseases, omitting rare events is also an important gap. In his *Evolutionary Biology of Parasites*, Price noted ‘The ecology of rare events, an important aspect of life for parasites, is yet to be developed’ (Price 1980). One such rare event of importance in EEID is host range shifts, whose rarity as a subject of study among the studies we reviewed is notable given their importance to public and animal health. Host shift events are the subject of intense study, much of which focuses on host shifts that have happened and the development of methods to predict when and where they will occur next (e.g., Becker et al. 2019 and papers referenced therein). However, the relative rarity with which host shifts occur makes them difficult to observe (at least in an unbiased way) in nature. As such, host shifts must be the object of experimental, model systems research. Such research can complement retrospective research on host shifts, by providing quantitative information

about the probability of unobserved events, such as the failure of a pathogen to spillover, and mechanistic explanations for spillover (and its absence) (Parrish et al. 2008; Gostic et al. 2019; Mollentze et al. 2020). Some models will be better than others for studying rare events like host shifts and other rare events, such as the addition/loss of host species as hosts or of transmission modes. We should prioritize systems where particularly vast sample sizes can be generated e.g. *Drosophila* viruses (Longdon et al. 2011, 2015) and where the pathogen can be safely contained (see discussions related to ‘gain of function’ experiments (Duprex et al. 2015)).

Toward Horizontal Integration or the comparative study of EEID

While our review was not designed to uncover comparative research, it is clear from reading the studies that we often do not perform research with a comparative mindset or with reference to an infrastructure that would facilitate such an approach to research.

Yet comparative studies could help us draw general lessons about the ecology and evolution of infectious diseases, as a number of recent studies have demonstrated (Leggett et al. 2012; Acevedo et al. 2019), and our review shows that there is ample room to use experimental data in this manner, at least as regards questions of (the interactions between) individual level disease processes (Fig. 3a). The systems reviewed herein share several features, merely by dint of being experimental systems used for EEID research. In all systems, infections can be generated, a metric of disease measured, and pathogens enumerated. In almost all systems, moreover, ‘individual’ level variables such as host strain, parasite strain, and parasite inoculum could be (or have been) simultaneously varied. As such, we are well-placed to ask key questions in EEID from a comparative perspective — e.g., are there thresholds for the establishment of infection and disease? How does pathogen burden vary with disease through time? Does disease predict transmission?

To systematically use model systems in this way, we need i) standardized experimental protocols, and ii) the infrastructure for sharing/storing both protocols, data generated and resources produced, such as parasite and host strains.

The project of designing standard experiments should be performed with reference to theory (e.g., Day 2002) and may itself stimulate theoretical research. Take the example of dose-response experiments, which are routinely conducted as part of system optimization but can also illuminate host-parasite interactions within-host. Should inoculum size be varied by a consistent scale (arithmetically, logarithmically) across systems and how should this change with pathogen type (e.g., bacteria vs. helminths, or macro vs. microparasite)? As of now, parasite inoculum is commonly varied logarithmically but there are other possibilities – e.g., might we consider varying dose with reference to relative biomass or metabolic rate, in the interests of understanding the energetic requirements of maintaining an infection? Theory can also help us with the prospective interpretation of data that has already been collected. To stay with the dose-response example, many dose-response experiments already exist (P.A. Clay, M.H. Cortez, and M.A. Duffy, *unpublished manuscript*) and there are likely many others that went unpublished. By illuminating how our inferences about disease processes are affected by study design, theory can enable these older datasets to be used to their fullest. Lastly, theory has a key role in providing precise definitions of key terms, without which we cannot hope to make comparisons between studies. Our review suggests common definitions are greatly needed since there was great variation in the way even the most commonly invoked and important concepts in EEID were defined. For example, in the papers we reviewed, virulence was defined in the same way that we have used it here, as well as to mean the harm an infection causes to the host (without respect to ‘who’ caused it), the capacity for a parasite to infect a host (Flor 1958; Barrett et al. 2007), and the capacity for a parasite to kill a host (Fenner 1949c).

In addition to standardized experimental designs, an infrastructure for the collation, storage, and sharing of protocols, data and resources could substantially advance comparative studies in EEID. The sharing of protocols and data is now easily facilitated by open source repositories such as protocols.io and datadryad.org. Yet the collation of such data into cohesive dataset is still an enormous challenge, albeit one with significant payoffs as the Global Mammal Parasite Database demonstrates (Stephens et al. 2017). Other collaborative efforts, such as NutNet, serve as instructive examples of how to systematize the entire process of research, from the design of experiments to data sharing, and the advantage of doing so (Adler et al. 2011; Fraser et al. 2013). In addition to sharing of experimental designs and data, sharing of organisms will also facilitate the expansion and replication of EEID research. Across the biological sciences, strain banks enable scientists to readily obtain specific strains of an organism and indeed, *P. chabaudi* and *H. polygyrus* can be thus acquired. However, the studies reviewed herein often use pathogen/host strains that are neither standardized (e.g. experimental infections are often generated using ‘isolates’ which can contain multiple pathogen genotypes (Luijckx et al. 2010)) nor broadly sharable. The use of polyclonal inocula or a diversity of hosts is often entirely appropriate for the questions at hand. But, where possible, the collection, preservation and maintenance of excess source organisms would enable post-hoc isolation of genetically-identical strains for comparative work. Establishing infrastructure for the sharing of data and resources will help us to replicate experiments, widen and democratize access to research resources, and, most relevant to the theme of this paper, generate general insights about disease processes.

Part III: Developing criteria for model systems in EEID

In parts I and II we used a concept of model systems borrowed from the biomolecular sciences as a reference point for a discussion of what ecologists and evolutionary biologists of infectious

disease use ‘model systems’ for and how they might use them better. But EEID has a different outlook and different priorities than the cellular-molecular-developmental fields of biology. Right off the bat, it was clear that this biomolecular conception of model systems was not a perfect fit. Model organisms in genetics, for example, are ‘not primarily studied because they are interesting in their own right’ (Ankeny and Leonelli 2011, pg. 318). By contrast, many of the model systems in EEID were established and are studied precisely because they are fascinating examples of evolution’s mischief or because they are of medical, agricultural, or conservation concern. As such, experimental systems in EEID serve *both* as interesting foci and are used to elucidate general rules about infectious diseases. This illustrates that EEID as a field may want to establish a bespoke definition of what a model system is and what it can be used for. In this part of the paper, we highlight issues that other fields have had to confront in their discussions about model systems, as a jumping off point for our own.

The essential idea that underpins model systems research is that we can learn about the general principles of nature from organisms that have features that make them well suited to research, because these model organisms share key features with other, less-easily studied organisms (Krebs 1975). That is to say, model organisms are *representative* of other organisms. Holmes traces this idea to Aristotle’s observation that different organisms share consistent body plans (Holmes 1993) and, of course, we now know that such similar traits are often (though not always) the result of evolutionary conservatism (Jenner and Wills 2007).

As we think about the design and use of model systems in EEID, then, we must ask ourselves at what level of biological organization do diseases have the equivalent of a similar ‘body plan’? Is phylogenetic relatedness what unites pathosystems and, if so, across what level of taxonomy can we generalize — e.g., can one virus only be a useful model of viruses in its family or of viruses in general? Assessing the representativeness of model systems based on their

taxonomy alone seems too limited. After all, the models developed from IEV and measles have been modified to explain the dynamics of a variety of pathogens including e.g., bacterial infections like pertussis. It seems more important that our model systems represent the characters or key concepts of interest to our field (Travis 2006; Jenner and Wills 2007), an idea implicit in the theory-first way we that conducted this review. The work of Anderson & May suggests that systems might be usefully grouped for comparative purposes into the categories of microparasite vs. macroparasite (Anderson and May 1991). Indeed, in performing this review, it felt ‘unnatural’ to lump together the replication of microparasites and maturation of macroparasites into one category (Table 2), since these processes involve different energetic requirements on the part of host and parasite, stimulate different sorts of immune defenses, and thus very likely have different evolutionary consequences. An alternative way to group pathosystems might be at the level of parasite life cycle: e.g., complex vs. simple, obligate- vs. non-obligate killer, which can greatly impact ecological and evolutionary dynamics (O’Keefe and Antonovics 2002). Importantly, it may not be enough to consider the traits of just parasites. EEID, unlike anatomy, genetics, or evolutionary developmental biology, requires model systems that represent the *relationship* between two organisms. We may thus need to pay attention to or define metrics that measure the interaction between pathogen and host — e.g., a measure of the *relative* speed of pathogen and host evolutionary rates, to elucidate which systems are representative of one another.

Much of the recent debate about the representativeness of model systems has focused on the extent to which the traits that make models easy to study also make them unrepresentative of organisms at large, and so bias the picture they give us of the natural world (Alfred and Baldwin 2015). For example, in his classic paper on what makes a model system, Krebs noted that one thing that makes model systems useful is the ‘magnitude of the phenomenon to be studied’ — i.e., that the model systems possess the trait of interest in abundance (Krebs 1975). Recently it

has been argued that organisms possessing extreme characters may be very derived, originating in relatively unusual unconstrained ecological and evolutionary contexts, and so are ill representative of (the processes that generate) biodiversity at large (Alfred and Baldwin 2015). This idea has driven a push toward extending the taxonomic sampling in genetics and other fields.

The extent to which the model systems reviewed here are biased deserves debate, as will the biases generated by any future-defined selection criteria for model systems. Ironically, the lack of a well-defined notion of a model system may have spared EEID some of the problems that arise due to the biased sampling of systems in other fields whereby they only exhibit the very extremes in the traits of interest or a specific taxonomic group. The majority of systems reviewed here were first studied either in another disease-focused field, like parasitology or immunology, as models of a specific disease, and/or because of their applied importance (Table 1). Thus, while they are united in being pathogenic, they represent a relatively random taxonomic set and, unlike many infections studied in the biomedical sciences, do not represent the most pathogenic of diseases. This set of pathosystems may not represent infectious diseases in other ways, however, most obviously because traits that make them easy to study in the laboratory may not be widely distributed in nature. For example, as we have noted, in many of the systems reviewed the (most commonly studied) host cannot fully recover. This feature makes these pathosystems easy to study — they can be modelled using a simple SI framework and, in the context of laboratory studies of epidemics, reduce the time required for sampling, since researchers only need to track individuals until infection is confirmed.

Are these systems thus unrepresentative of the many infections that do resolve? In some ways, certainly. But that does not mean they entirely uninformative about the process of recovery. Take an analogy from evolutionary developmental biology. Jenner & Wills (2007) countered the

argument that systems invulnerable to environmental change were poorly placed for understanding environmental influence on development, by noting that such systems could help evolutionary developmental biologists understand the complementary phenomenon of canalization. In a similar vein, non-resolving diseases might help illuminate much about tolerance, a host response complementary to, and which coexists alongside, resistance.

Indeed, it might be argued that the ways in which model systems fail to represent other systems are just as, if not more, useful than the ways in which they are. Take the example of *M. lini* and *P. ramosa*. Interactions of these two pathogens and their hosts are mediated by ‘matching’ interactions between their genes and that of their hosts. These systems are often used by EEID researchers to ask similar questions, for example the impact of space and metapopulations on disease dynamics. Yet, their coevolutionary dynamics are different because of the details of what happens when a parasite and host genes ‘match’ (mediating ‘incompatibility’ and ‘compatibility’, respectively) (Thrall et al. 2016). So, details matter, and teach us where our theory falls down and motivate us to make it anew. This fact is not at odds with the program of model systems research, however, for we cannot understand whether the details are consistent with, or contrary to, theory if we do not set out to do research in a comparative context.

Conclusion

As a relatively young field, EEID nonetheless has a number of systems that are used as touchstones. While the earliest systems were laboratory-based and borrowed from other fields, with time we have developed our own. These newer systems, which often incorporate the field and lab, give us access to greater variation in hosts and parasites and permit the exploration of processes at different scales. Yet success has been variable. Laboratory models have been used with limited success to address foundational questions about the effect of parasites on population or ecosystem level processes and vice versa. A concerted effort toward vertical

integration within systems would address this gap. Similarly, systematic horizontal integration between systems could help us to establish whether phenomena observed in one system can be generalized to others and to elucidate general rules underlying disease transmission. Indeed, as we have learned during the writing of this piece, the very exercise of asking ‘what do we want from our model systems?’ could help to establish a concrete, consensus definitions of concepts or experimental designs that could be used across the model systems in EEID.

Acknowledgments

We would like to thank Janis Antonovics, Jukka Jokela, and Andrew Read for providing helpful insights about the changing role of model systems through EEID’s history, Aaron King for providing helpful comments on an earlier draft of the manuscript and two anonymous reviewers and the editor for comments that greatly improved the manuscript. This work was supported by the National Science Foundation (NSF-1655856) and by the Moore Foundation (GBMF9202; DOI: <https://doi.org/10.37807/GBMF9202>).

Author contributions

NW conceived of the framing of the paper, performed the literature review, and wrote the paper. MAD provided feedback on the framing of the paper and edited the manuscript.

References

- Acevedo, M. A., F. P. Dilleuth, A. J. Flick, M. J. Faldyn, and B. D. Elderd. 2019. Virulence-driven trade-offs in disease transmission: A meta-analysis*. *Evolution* 73:636–647.
- Achtman, A., R. Stephens, E. Cadman, V. Harrison, and J. Langhorne. 2007. Malaria-specific antibody responses and parasite persistence after infection of mice with *Plasmodium chabaudi chabaudi*. *Parasite Immunology* 29:435–444.
- Adelman, J. S., S. C. Moyers, D. R. Farine, and D. M. Hawley. 2015. Feeder use predicts both acquisition and transmission of a contagious pathogen in a North American songbird. *Proceedings of the Royal Society B: Biological Sciences* 282:20151429.
- Adler, P. B., E. W. Seabloom, E. T. Borer, H. Hillebrand, Y. Hautier, A. Hector, W. S. Harpole, et al. 2011. Productivity Is a Poor Predictor of Plant Species Richness. *Science* 333:1750–1753.
- Alexander, H. M. 1989. An experimental field study of anther-smut disease of *Silene alba* caused by *Ustilago violacea*: genotypic variation and disease incidence. *Evolution* 43:835–847.
- Alfred, J., and I. T. Baldwin. 2015. New opportunities at the wild frontier. *eLife* 4:e06956.
- Anderson, R. M., and R. M. May. 1979. Population biology of infectious diseases: Part I. *Nature* 280:361–367.
- Anderson, R. M., and R. M. May. 1991. A framework for discussing the population biology of infectious diseases. Pages 1–7 *in* *Infectious Diseases of Humans: Dynamics and Control*.
- Ankeny, R. A., and S. Leonelli. 2011. What's so special about model organisms? *Studies in History and Philosophy of Science Part A* 42:313–323.
- Antonovics, J., M. Hood, and J. Partain. 2002. The ecology and genetics of a host shift: *Microbotryum* as a model system. *The American Naturalist* 160 Suppl 4:S40-53.
- Babayan, S. A., R. J. Orton, and D. G. Streicker. 2018. Predicting reservoir hosts and arthropod vectors from evolutionary signatures in RNA virus genomes. *Science* 362:577–580.
- Bailes, E. J., J. Bagi, J. Coltman, M. T. Fountain, L. Wilfert, and M. J. F. Brown. 2020. Host density drives viral, but not trypanosome, transmission in a key pollinator. *Proceedings of the Royal Society B* 287:20191969.

- Bansemir, A. D., and M. V. K. Sukhdeo. 1996. Villus length influences habitat selection by *Heligmosomoides polygyrus*. *Parasitology* 113:311–316.
- Barber, I., and J. P. Scharsack. 2010. The three-spined stickleback-*Schistocephalus solidus* system: an experimental model for investigating host-parasite interactions in fish. *Parasitology* 137:411–424.
- Barrett, L. G., P. H. Thrall, and J. J. Burdon. 2007. Evolutionary diversification through hybridization in a wild host-pathogen interaction. *Evolution* 61:1613–1621.
- Becker, D. J., A. D. Washburne, C. L. Faust, J. R. C. Pulliam, E. A. Mordecai, J. O. Lloyd-Smith, and R. K. Plowright. 2019. Dynamic and integrative approaches to understanding pathogen spillover. *Philosophical Transactions of the Royal Society B: Biological Sciences* 374:20190014.
- Bell, A. S., S. Huijben, K. P. Paaijmans, D. G. Sim, B. H. K. Chan, W. A. Nelson, and A. F. Read. 2012. Enhanced transmission of drug-resistant parasites to mosquitoes following drug treatment in rodent malaria. *PLoS ONE* 7:e37172.
- Benmayor, R., D. J. Hodgson, G. G. Perron, and A. Buckling. 2009. Host Mixing and Disease Emergence. *Current biology* : CB 19:764–767.
- Betterton, C. 1974. Studies on the host specificity of the eyefluke, *Diplostomum spathaceum*, in brown and rainbow trout. *Parasitology* 69:11–29.
- Birger, R. B., R. D. Kouyos, T. Cohen, and E. C. Griffiths. 2015. The potential impact of coinfection on antimicrobial chemotherapy and drug resistance. *Trends in Microbiology*.
- Bonneaud, C., M. Giraudeau, L. Tardy, M. Staley, G. E. Hill, and K. J. McGraw. 2018. Rapid Antagonistic Coevolution in an Emerging Pathogen and Its Vertebrate Host. *Current Biology* 28:2978-2983.e5.
- Bonneaud, C., L. Tardy, M. Giraudeau, G. E. Hill, K. J. McGraw, and A. J. Wilson. 2019. Evolution of both host resistance and tolerance to an emerging bacterial pathogen. *Evolution Letters* 3:544–554.
- Bouwman, K. M., and D. M. Hawley. 2010. Sickness behaviour acting as an evolutionary trap? Male house finches preferentially feed near diseased conspecifics. *Biology Letters* 6:462–465.
- Brandell, E. E., D. J. Becker, L. Sampso, and K. M. Forbes. 2020. The rise of disease ecology.
- Brindley, P. J., and C. Dobson. 1981. Genetic control of liability to infection with *Nematospiroides dubius* in mice: selection of refractory and liable populations of mice. *Parasitology* 83:51–65.
- Buckling, A. G. J., L. H. Taylor, J. M. –R Carlton, and A. F. Read. 1997. Adaptive changes in *Plasmodium* transmission strategies following chloroquine chemotherapy. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 264:553–559.

- Budischak, S. A., K. Sakamoto, L. C. Megow, K. R. Cummings, J. F. U. Jr, and V. O. Ezenwa. 2015. Resource limitation alters the consequences of co-infection for both hosts and parasites. *International journal for parasitology* 45:455–463.
- Buechel, S. D., and P. Schmid-Hempel. 2016. Colony pace: a life-history trait affecting social insect epidemiology. *Proceedings of the Royal Society B: Biological Sciences* 283:20151919.
- Carpenter, S. R. 1996. Microcosm experiments have limited relevance for community and ecosystem ecology. *Ecology* 77:677–680.
- Chandler, A. 1939. The effects of number and age of worms on development of primary and secondary infections with *Hymneolepis diminuta* in rats, and an investigation into the true nature of “premunitio” in tapeworm infections. *American Journal of Hygiene* 105–114.
- Chao, L., B. R. Levin, and F. M. Stewart. 1977. A Complex Community in a Simple Habitat: An Experimental Study with Bacteria and Phage. *Ecology* 58:369–378.
- Chappell, L. H., and A. W. Pike. 1976. Loss of *Hymenolepis diminuta* from the rat. *International Journal for Parasitology* 6:333–339.
- Chaudhri, G., V. Panchanathan, R. M. L. Buller, A. J. M. van den Eertwegh, E. Claassen, J. Zhou, R. de Chazal, et al. 2004. Polarized type 1 cytokine response and cell-mediated immunity determine genetic resistance to mousepox. *Proceedings of the National Academy of Sciences* 101:9057–9062.
- Choo, K., P. D. Williams, and T. Day. 2003. Host mortality, predation and the evolution of parasite virulence. *Ecology Letters* 6:310–315.
- Clay, P. A., M. H. Cortez, M. A. Duffy, and V. H. W. Rudolf. 2019. Priority effects within coinfecting hosts can drive unexpected population-scale patterns of parasite prevalence. *Oikos* 128:571–583.
- Clerc, M., D. Ebert, and M. D. Hall. 2015. Expression of parasite genetic variation changes over the course of infection: implications of within-host dynamics for the evolution of virulence. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 282:20142820–20142820.
- Cressler, C. E., D. V. McLEOD, C. Rozins, J. van den Hoogen, and T. Day. 2016. The adaptive evolution of virulence: a review of theoretical predictions and empirical tests. *Parasitology* 143:915–930.
- Crowden, A. E., and D. M. Broom. 1980. Effects of the eyefluke, *Diplostomum spathaceum* on the behaviour of dace (*Leuciscus leuciscus*). *Animal Behaviour* 28:287–294.
- Day, T. 2002. On the evolution of virulence and the relationship between various measures of mortality. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 269:1317–1323.

- Day, T., S. Alizon, and N. Mideo. 2011. Bridging scales in the evolution of infectious disease life histories: theory. *Evolution* 65:3448–3461.
- deRoode, J., R. Pansini, S. Cheesman, M. Helsinki, S. Huijben, A. Wargo, A. Bell, et al. 2005. Virulence and competitive ability in genetically diverse malaria infections. *Proceedings of the National Academy of Sciences* 102:7624–7628.
- Dhondt, A. A., S. Altizer, E. G. Cooch, A. K. Davis, A. Dobson, M. J. L. Driscoll, B. K. Hartup, et al. 2005. Dynamics of a novel pathogen in an avian host: Mycoplasmal conjunctivitis in house finches. *Acta Tropica* 94:77–93.
- Dietrich, M. R., R. A. Ankeny, N. Crowe, S. Green, and S. Leonelli. 2019. How to choose your research organism. *Studies in History and Philosophy of Science Part C: Studies in History and Philosophy of Biological and Biomedical Sciences* 101227.
- Dobson, C., and M. E. Owen. 1977. Influence of serial passage on the infectivity and immunogenicity of *Nematospiroides dubius* in mice. *International Journal for Parasitology* 7:463–466.
- . 1978. Effect of host sex on passive immunity in mice infected with *Nematospiroides dubius*. *International Journal for Parasitology* 8:359–364.
- Duffy, M. A., C. E. Cáceres, and S. R. Hall. 2019. Healthy herds or predator spreaders? Insights from the plankton into how predators suppress and spread disease. Duffy, M., Cáceres, C., & Hall, S. (2019). Healthy herds or predator spreaders? Insights from the plankton into how predators suppress and spread disease. In K. Wilson, A. Fenton, & D. Tompkins (Eds.), *Wildlife Disease Ecology: Linking Theory to Data and Application* (Ecological Reviews, pp. 458-479). Cambridge: Cambridge University Press. doi:10.1017/9781316479964.016 458–479.
- Duncan, A. B., and T. J. Little. 2007. Parasite-driven genetic change in a natural population of *Daphnia*. *Evolution* 61:796–803.
- Duneau, D., P. Luijckx, F. Ben-Ami, C. Laforsch, and D. Ebert. 2011. Resolving the infection process reveals striking differences in the contribution of environment, genetics and phylogeny to host-parasite interactions. *BMC biology* 9:11.
- Duprex, W. P., R. A. M. Fouchier, M. J. Imperiale, M. Lipsitch, and D. A. Relman. 2015. Gain-of-function experiments: time for a real debate. *Nature Reviews Microbiology* 13:58–64.
- Dybdahl, M. F., and C. M. Lively. 1998. Host-parasite coevolution: evidence for rare advantage and time-lagged selection in a natural population. *Evolution* 52:1057–1066.
- Ebert, D. 2008. Host–parasite coevolution: Insights from the *Daphnia*–parasite model system. *Current Opinion in Microbiology* 11:290–301.
- Ebert, D., M. Lipsitch, and K. L. Mangin. 2000. The effect of parasites on host population density and extinction: Experimental epidemiology with *Daphnia* and six microparasites. *The American Naturalist* 156:459–477.

Ezenwa, V. O., and A. E. Jolles. 2015. Opposite effects of anthelmintic treatment on microbial infection at individual versus population scales. *Science*.

Fenner, F. 1947a. Studies in infectious ectromelia of mice. I. Immunization of mice against ectromelia with living vaccinia virus. *Australian Journal of Experimental Biology and Medical Science* 25:257–274.

———. 1947b. Studies in infectious ectromelia in mice II. Natural transmission: the portal of entry of the virus. *Australian Journal of Experimental Biology and Medical Science* 275–282.

———. 1948a. The epizootic behaviour of mousepox (infectious ectromelia of mice) II. The course of events in long-continued epidemics. *Australian Journal of Experimental Biology and Medical Science*, *Australian Journal of Experimental Biology and Medical Science* 46:383–393.

———. 1948b. The epizootic behaviour of mouse-pox, infectious ectromelia. *British journal of experimental pathology* 29:69–91.

———. 1949a. Studies in mousepox (infectious ectromelia of mice). IV. Quantitative investigations on the spread of virus through the host in actively and passively immunized animals. *Australian Journal of Experimental Biology and Medical Science* 27:1–18.

———. 1949b. Studies in infectious ectromelia in mice. VII. The effect of the age of the host upon the response to infection. *Australian Journal of Experimental Biology and Medical Science* 27:45–53.

———. 1949c. Studies in mousepox (infectious ectromelia of mice). VI. A comparison of the virulence and infectivity of three strains of ectromelia virus. *Australian Journal of Experimental Biology and Medical Science* 27:31–44.

———. 2000. Adventures with poxviruses of vertebrates. *FEMS Microbiology Reviews* 24:123–133.

Fenton, A., D. G. Streicker, O. L. Petchey, and A. B. Pedersen. 2015. Are all hosts created equal? Partitioning host species contributions to parasite persistence in multihost communities. *The American Naturalist*.

Ferguson, H. M., and A. F. Read. 2002. Genetic and environmental determinants of malaria parasite virulence in mosquitoes. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 269:1217–1224.

Flor, H. 1971. Current status of the gene-for-gene concept. *Annual Reviews in Phytopathology* 9:275–296.

Flor, H. H. 1958. Mutation to wider virulence in *Melampsora lini*. *Phytopathology* 48:297–301.

———. 1960. The inheritance of X-Ray-induced mutations to virulence in a urediospore culture of Race-1 of *Melampsora lini*. *Phytopathology* 603–605.

- Fouks, B., E. G. Robb, and H. M. G. Lattorff. 2018. Role of conspecifics and personal experience on behavioral avoidance of contaminated flowers by bumblebees. *Current Zoology* 65:447–455.
- Fraser, L. H., H. A. Henry, C. N. Carlyle, S. R. White, C. Beierkuhnlein, J. F. Cahill, B. B. Casper, et al. 2013. Coordinated distributed experiments: an emerging tool for testing global hypotheses in ecology and environmental science. *Frontiers in Ecology and the Environment* 11:147–155.
- Gegear, R. J., M. C. Otterstatter, and J. D. Thomson. 2006. Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proceedings of the Royal Society B: Biological Sciences* 273:1073–1078.
- Gipson, S. A. Y., L. Jimenez, and M. D. Hall. 2019. Host sexual dimorphism affects the outcome of within-host pathogen competition. *Evolution* 73:1443–1455.
- Gonzalez, A., J. C. Clemente, A. Shade, J. L. Metcalf, S. Song, B. Prithviraj, B. E. Palmer, et al. 2011. Our microbial selves: what ecology can teach us. *EMBO reports* 12:775–784.
- Goodman, B. A., and P. T. J. Johnson. 2011. Disease and the extended phenotype: Parasites control host performance and survival through induced changes in body plan. *PLoS ONE* 6:e20193.
- Gostic, K. M., E. A. Wunder, V. Bisht, C. Hamond, T. R. Julian, A. I. Ko, and J. O. Lloyd-Smith. 2019. Mechanistic dose–response modelling of animal challenge data shows that intact skin is a crucial barrier to leptospiral infection. *Philosophical Transactions of the Royal Society B: Biological Sciences* 374:20190367.
- Graham, A. L., T. J. Lamb, A. F. Read, and J. E. Allen. 2005. Malaria-Filaria Coinfection in Mice Makes Malarial Disease More Severe unless Filarial Infection Achieves Patency. *The Journal of Infectious Diseases* 191:410–421.
- Graham, A. L., D. M. Shuker, L. C. Pollitt, S. K. J. R. Auld, A. J. Wilson, and T. J. Little. 2010. Fitness consequences of immune responses: strengthening the empirical framework for ecoimmunology. *Functional Ecology* 25:5–17.
- Gregory, R. D., A. E. Keymer, and J. R. Clarke. 1990. Genetics, sex and exposure: The ecology of *Heligmosomoides polygyrus* (Nematoda) in the wood mouse. *The Journal of Animal Ecology* 59:363.
- Hakalahti, T., A. Karvonen, and E. T. Valtonen. 2006. Climate warming and disease risks in temperate regions – *Argulus coregoni* and *Diplostomum spathaceum* as case studies. *Journal of Helminthology* 80:93–98.
- Hall, M. D., and D. Ebert. 2012. Disentangling the influence of parasite genotype, host genotype and maternal environment on different stages of bacterial infection in *Daphnia magna*. *Proceedings of the Royal Society B: Biological Sciences* 279:3176–3183.

- Hansen, J., and T. Day. 2014. Coinfection and the evolution of drug resistance. *Journal of evolutionary biology* 27:2595–2604.
- Hawley, D. M., E. E. Osnas, A. P. Dobson, W. M. Hochachka, D. H. Ley, and A. A. Dhondt. 2013. Parallel patterns of increased virulence in a recently emerged wildlife pathogen. *PLoS Biology* 11:e1001570.
- Hawley, D. M., K. V. Sydenstricker, G. V. Kollias, and A. A. Dhondt. 2005. Genetic diversity predicts pathogen resistance and cell-mediated immunocompetence in house finches. *Biology Letters* 1:326–329.
- Hernandez, A. D., and M. V. Sukhdeo. 1995. Host grooming and the transmission strategy of *Heligmosomoides polygyrus*. *The Journal of Parasitology* 81:865–9.
- Hite, J. L., A. C. Pfenning, and C. E. Cressler. 2019. Starving the Enemy? Feeding Behavior Shapes Host-Parasite Interactions. *Trends in Ecology & Evolution* 35:68–80.
- Holmes, F. L. 1993. The old martyr of science: The frog in experimental physiology. *Journal of the History of Biology* 26:311–328.
- Holmes, J. C. 1959. Competition for carbohydrate between the rat tapeworm, *Hymenolepis diminuta*, and the Acanthocephalan, *Moniliformis dubius*. Page 31 in Program and Abstracts of the Thirty-Fourth Annual Meeting of the American Society of Parasitologists (Vol. 45). Presented at the 34th Annual Meeting of the American Society of Parasitologists, *The Journal of Parasitologists*.
- Horne, M. T. 1970. Coevolution of *Escherichia coli* and Bacteriophages in Chemostat Culture. *Science* 168:992–993.
- Hudson, P. J., and D. Newborn. 1998. Prevention of population cycles by parasite removal. *Science* 282:2256–2258.
- Hughes, D. 2013. Pathways to understanding the extended phenotype of parasites in their hosts. *Journal of Experimental Biology* 216:142–147.
- Huijben, S., B. H. K. Chan, W. A. Nelson, and A. F. Read. 2018. The impact of within-host ecology on the fitness of a drug resistant parasite. *Evolution, Medicine, and Public Health* 2018:eoy016-.
- Huijben, S., B. H. K. Chan, and A. F. Read. 2015. Relevance of undetectably rare resistant malaria parasites in treatment failure: experimental evidence from *Plasmodium chabaudi*. *The American Journal of Tropical Medicine and Hygiene* 92:1214–1221.
- Jenner, R. A., and M. A. Wills. 2007. The choice of model organisms in evo–devo. *Nature Reviews Genetics* 8:311–314.
- Jent, D., A. Perry, J. Critchlow, and A. T. Tate. 2019. Natural variation in the contribution of microbial density to inducible immune dynamics. *Molecular Ecology* 28:5360–5372.

Johnson, P. T. J., D. M. Calhoun, T. B. Riepe, and J. Koprivnikar. 2019. Chance or choice? Understanding parasite selection and infection in multi-host communities. *International Journal for Parasitology* 49:407–415.

Johnson, P. T. J., J. M. Chase, K. L. Dosch, R. B. Hartson, J. A. Gross, D. J. Larson, D. R. Sutherland, et al. 2007. Aquatic eutrophication promotes pathogenic infection in amphibians. *Proceedings of the National Academy of Sciences* 104:15781–15786.

Johnson, P. T. J., D. L. Preston, J. T. Hoverman, and K. L. D. Richgels. 2014. Biodiversity decreases disease through predictable changes in host community competence. *Nature* 494:230–233.

Johnson, P. T. J., J. R. Rohr, J. T. Hoverman, E. Kellermanns, J. Bowerman, and K. B. Lunde. 2012. Living fast and dying of infection: host life history drives interspecific variation in infection and disease risk. *Ecology Letters* 15:235–242.

Johnson, P. T. J., D. R. Sutherland, J. M. Kinsella, and K. B. Lunde. 2004. Review of the Trematode Genus *Ribeiroia* (Psilostomidae): Ecology, Life History and Pathogenesis with Special Emphasis on the Amphibian Malformation Problem. *Advances in Parasitology*, *Advances in Parasitology* 57:191–253.

Johnson, P. T., K. B. Lunde, R. W. Haight, J. Bowerman, and A. R. Blaustein. 2001. *Ribeiroia ondatrae* (Trematoda: Digenea) infection induces severe limb malformations in western toads (*Bufo boreas*). *Canadian Journal of Zoology* 79:370–379.

Johnson, P. T., D. L. Preston, J. T. Hoverman, and B. E. LaFonte. 2013. Host and parasite diversity jointly control disease risk in complex communities. *Proceedings of the National Academy of Sciences of the United States of America* 110:16916–16921.

Jones, K. E., N. G. Patel, M. a Levy, A. Storeygard, D. Balk, J. L. Gittleman, and P. Daszak. 2008. Global trends in emerging infectious diseases. *Nature* 451:990–993.

Karvonen, A., S. Paukuu, E. T. Valtonen, and P. Hudson. 2003. Transmission, infectivity and survival of *Diplostomum spathaceum* cercariae. *Parasitology* 127:217–224.

Karvonen, A., O. Seppala, and E. T. Valtonen. 2004. Parasite resistance and avoidance behaviour in preventing eye fluke infections in fish. *Parasitology* 129:159–164.

Katz, P. S. 2016. 'Model organisms' in the light of evolution. *Current Biology* 26:R649–R650.

Kavaliers, M., D. D. Colwell, W. J. Braun, and E. Choleris. 2003. Brief exposure to the odour of a parasitized male alters the subsequent mate odour responses of female mice. *Animal Behaviour* 65:59–68.

Keeling, M. J., and P. Rohani. 2011. *Modeling Infectious Diseases in Humans and Animals*. Princeton University Press. Princeton University Press.

Keesing, F., L. K. Belden, P. Daszak, A. Dobson, C. D. Harvell, R. D. Holt, P. Hudson, et al. 2010. Impacts of biodiversity on the emergence and transmission of infectious diseases. *Nature* 468:647–652.

Keymer, A. E. 1985. Experimental epidemiology: *Nematosporoides dubius* and the laboratory mouse. Pages 55–76 in D. Rollinson and R. Anderson, eds. *Ecology and Genetics of Host-Parasite Interactions*, Linnean Society Symposium Series (Vol. 11). Academic Press.

Keymer, A. E., and R. M. Anderson. 1979. The dynamics of infection of *Tribolium confusum* by *Hymenolepis diminuta*: the influence of infective-stage density and spatial distribution. *Parasitology* 79:195–207.

Klemme, I., and A. Karvonen. 2018. Shoaling with infected conspecifics does not improve resistance to trematode infection. *Ethology* 124:170–176.

Knowles, S. C. L., A. Fenton, O. L. Petchey, T. R. Jones, R. Barber, and A. B. Pedersen. 2013. Stability of within-host-parasite communities in a wild mammal system. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 280:20130598–20130598.

Koch, H., and P. Schmid-Hempel. 2011. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proceedings of the National Academy of Sciences* 108:19288–19292.

Koch, H., and P. Schmid-Hempel. 2012. Gut microbiota instead of host genotype drive the specificity in the interaction of a natural host-parasite system. *Ecology Letters* 15:1095–1103.

Kollias, G. V., K. V. Sydenstricker, H. W. Kollias, D. H. Ley, P. R. Hosseini, V. Connolly, and A. A. Dhondt. 2004. Experimental infection of house finches with *Mycoplasma gallisepticum*. *Journal of Wildlife Diseases* 40:79–86.

Koskella, B. 2014. Bacteria-Phage Interactions across Time and Space: Merging Local Adaptation and Time-Shift Experiments to Understand Phage Evolution*. *The American Naturalist* 184:S9–S21.

Koskella, B., L. J. Hall, and C. J. E. Metcalf. 2017. The microbiome beyond the horizon of ecological and evolutionary theory. *Nature Ecology & Evolution* 1:1606–1615.

Krebs, H. A. 1975. The August Krogh principle: “For many problems there is an animal on which it can be most conveniently studied.” *Journal of Experimental Zoology* 194:221–226.

Lafferty, K. D. 2009. The ecology of climate change and infectious diseases. *Ecology* 90:888–900.

LaFonte, B. E., and P. T. J. Johnson. 2013. Experimental infection dynamics: using immunosuppression and *in vivo* parasite tracking to understand host resistance in an amphibian–trematode system. *The Journal of Experimental Biology* 216:3700–3708.

- Lass, S., P. J. Hudson, J. Thakar, J. Saric, E. Harvill, R. Albert, and S. E. Perkins. 2013. Generating super-shedders: co-infection increases bacterial load and egg production of a gastrointestinal helminth. *Journal of the Royal Society Interface* 10:20120588.
- Leggett, H. C., C. K. Cornwallis, and S. a West. 2012. Mechanisms of Pathogenesis, Infective Dose and Virulence in Human Parasites. *PLoS Pathogens* 8:e1002512-5.
- Lenski, R. E., and B. R. Levin. 1985. Constraints on the Coevolution of Bacteria and Virulent Phage: A Model, Some Experiments, and Predictions for Natural Communities. *The American Naturalist* 125:585–602.
- Leonelli, S., and R. A. Ankeny. 2013. What makes a model organism? *Endeavour* 37:209–12.
- Lepak, J., and V. Thatcher. 1962. The development of *Nematospiroides dubius* in an abnormal site in the white mouse and some apparent effects on the serum proteins of the host. *Texas Reports on Biology and Medicine* 20:374–83.
- Lin, J.-D., J. C. Devlin, F. Yeung, C. McCauley, J. M. Leung, Y.-H. Chen, A. Cronkite, et al. 2020. Rewilding Nod2 and Atg16l1 Mutant Mice Uncovers Genetic and Environmental Contributions to Microbial Responses and Immune Cell Composition. *Cell Host & Microbe* 27:830-840.e4.
- Little, T. J., D. M. Shuker, N. Colegrave, T. Day, and A. L. Graham. 2010. The Coevolution of Virulence: Tolerance in Perspective. *PLoS Pathogens* 6:e1001006.
- Lively, C. M., and M. F. Dybdahl. 2000. Parasite adaptation to locally common host genotypes. *Nature* 405:679–681.
- Lively, C. M., J. C. de Roode, M. A. Duffy, A. L. Graham, and B. Koskella. 2014. Interesting Open Questions in Disease Ecology and Evolution*. *The American Naturalist* 184:S1–S8.
- Lloyd-Smith, J. O., D. George, K. M. Pepin, V. E. Pitzer, J. R. C. Pulliam, A. P. Dobson, P. J. Hudson, et al. 2009. Epidemic Dynamics at the Human-Animal Interface. *Science* 326:1362–1367.
- Longdon, B., J. D. Hadfield, J. P. Day, S. C. L. Smith, J. E. McGonigle, R. Cogni, C. Cao, et al. 2015. The Causes and Consequences of Changes in Virulence following Pathogen Host Shifts. *PLoS Pathogens* 11:e1004728.
- Longdon, B., L. Wilfert, J. Osei-Poku, H. Cagney, D. J. Obbard, and F. M. Jiggins. 2011. Host-switching by a vertically transmitted rhabdovirus in *Drosophila*. *Biology letters* 7:747–750.
- Luijckx, P., F. Ben-Ami, L. Mouton, L. D. Pasquier, and D. Ebert. 2010. Cloning of the unculturable parasite *Pasteuria ramosa* and its *Daphnia* host reveals extreme genotype-genotype interactions. *Ecology letters* 14:125–31.
- Marchetto, K. M., and A. G. Power. 2018. Coinfection Timing Drives Host Population Dynamics through Changes in Virulence. *The American Naturalist* 191:173–183.

- May, R. M., and R. M. Anderson. 1979. Population biology of infectious diseases: Part II. *Nature* 280:455–461.
- McKee, K. M., J. Koprivnikar, P. T. J. Johnson, and M. T. Arts. 2020. Parasite infectious stages provide essential fatty acids and lipid-rich resources to freshwater consumers. *Oecologia* 192:477–488.
- Mideo, N., A. Acosta-Serrano, T. Aebischer, M. J. F. Brown, A. Fenton, V.-P. Friman, O. Restif, et al. 2013a. Life in cells, hosts, and vectors: Parasite evolution across scales. *Infection, genetics and evolution* 13:344–347.
- . 2013b. Life in cells, hosts, and vectors: Parasite evolution across scales. *Infection, genetics and evolution* 13:344–347.
- Mideo, N., W. A. Nelson, S. E. Reece, A. S. Bell, A. F. Read, and T. Day. 2011. Bridging scales in the evolution of infectious disease life histories: application. *Evolution* 65:3298–3310.
- Mikheev, V. N., A. F. Pasternak, A. Y. Morozov, and J. Taskinen. 2019. Innate antipredator behavior can promote infection in fish even in the absence of predators. *Behavioral Ecology* 31:267–276.
- Milotic, D., M. Milotic, and J. Koprivnikar. 2017. Effects of road salt on larval amphibian susceptibility to parasitism through behavior and immunocompetence. *Aquatic Toxicology* 189:42–49.
- Mollentze, N., R. Biek, and D. G. Streicker. 2014. The role of viral evolution in rabies host shifts and emergence. *Current Opinion in Virology* 8:1–6.
- Mollentze, N., D. G. Streicker, P. R. Murcia, K. Hampson, and R. Biek. 2020. Virulence mismatches in index hosts shape the outcomes of cross-species transmission. *Proceedings of the National Academy of Sciences* 117:28859–28866.
- Mordecai, E. A., J. M. Cohen, M. V. Evans, P. Gudapati, L. R. Johnson, C. A. Lippi, K. Miazgowicz, et al. 2017. Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLoS Neglected Tropical Diseases* 11:e0005568.
- Morgan, M., J. Behnke, J. Lucas, and J. Peberdy. 1997. *In vitro* assessment of the influence of nutrition, temperature and larval density on trapping of the infective larvae of *Heligmosomoides polygyrus* by *Arthrobotrys oligospora*, *Duddingtonia flagrans* and *Monacrosporium megalosporum*. *Parasitology* 115:303–310.
- Morran, L. T., O. G. Schmidt, I. A. Gelarden, R. C. Parrish, and C. M. Lively. 2011. Running with the Red Queen: Host-Parasite Coevolution Selects for Biparental Sex. *Science* 333:216–218.
- Niewiadomska, K. 1986. Verification of the life-cycles of *Diplostomum spathaceum* (Rudolphi, 1819) and *D. pseudospathaceum* Niewiadomska, 1984 (Trematoda, Diplostomidae). *Systematic Parasitology* 8:23–31.

- O'Keefe, K. J., and J. Antonovics. 2002. Playing by different rules: the evolution of virulence in sterilizing pathogens. *The American Naturalist* 159:597–605.
- Olival, K. J., P. R. Hosseini, C. Zambrana-Torrel, N. Ross, T. L. Bogich, and P. Daszak. 2017. Host and viral traits predict zoonotic spillover from mammals. *Nature* 546:646–650.
- Orlowski, S. A., R. C. Jadin, D. L. Preston, and P. T. J. Johnson. 2012. Parasite transmission in complex communities: Predators and alternative hosts alter pathogenic infections in amphibians. *Ecology* 93:1247–1253.
- Otterstatter, M. C., and J. D. Thomson. 2007. Contact networks and transmission of an intestinal pathogen in bumble bee (*Bombus impatiens*) colonies. *Oecologia* 154:411–421.
- Overholt, E. P., M. A. Duffy, M. P. Meeks, T. H. Leach, and C. E. Williamson. 2020. Light exposure decreases infectivity of the *Daphnia* parasite *Pasteuria ramosa*. *Journal of Plankton Research* 42:41–44.
- Overholt, E. P., S. R. Hall, C. E. Williamson, C. K. Meikle, M. A. Duffy, and C. E. Cáceres. 2012. Solar radiation decreases parasitism in *Daphnia*. *Ecology Letters* 15:47–54.
- Packer, C., R. D. Holt, P. J. Hudson, K. D. Lafferty, and A. P. Dobson. 2003. Keeping the herds healthy and alert: implications of predator control for infectious disease. *Ecology Letters*, *Ecology Letters* 6:797–802.
- Palmer-Young, E. C., L. Ngor, R. B. Nevarez, J. A. Rothman, T. R. Raffel, and Q. S. McFrederick. 2019. Temperature dependence of parasitic infection and gut bacterial communities in bumble bees. *Environmental Microbiology* 21:4706–4723.
- Park, T. 1948. Interspecies competition in populations of *Trilobium confusum* Duval and *Trilobium castaneum* Herbst. *Ecological Monographs* 18:265–307.
- Parrish, C. R., E. C. Holmes, D. M. Morens, E. C. Park, D. S. Burke, C. H. Calisher, C. A. Laughlin, et al. 2008. Cross-Species Virus Transmission and the Emergence of New Epidemic Diseases. *Microbiology and Molecular Biology Reviews* 72:457–470.
- Paull, S. H., and P. T. Johnson. 2011. High temperature enhances host pathology in a snail–trematode system: possible consequences of climate change for the emergence of disease. *Freshwater Biology* 56:767–778.
- Ponton, F., F. Lalubin, C. Fromont, K. Wilson, C. Behm, and S. J. Simpson. 2011. Hosts use altered macronutrient intake to circumvent parasite-induced reduction in fecundity. *International Journal for Parasitology* 41:43–50.
- Preston, D. L., J. A. Mischler, A. R. Townsend, and P. T. J. Johnson. 2016. Disease Ecology Meets Ecosystem Science. *Ecosystems* 19:737–748.
- Price, P. W. 1980. *Evolutionary Biology of Parasites*. Princeton University Press. Princeton University Press.

- Raberg, L., D. Sim, and A. F. Read. 2007. Disentangling Genetic Variation for Resistance and Tolerance to Infectious Diseases in Animals. *Science* 318:812–814.
- Ravensdalle, M., A. Nemri, P. H. Thrall, J. Ellis, and P. Dodds. 2011. Co-evolutionary interactions between host resistance and pathogen effector genes in flax rust disease. *Molecular Plant Pathology* 12:93–102.
- Read, C. P. 1967. Longevity of the Tapeworm, *Hymenolepis diminuta*. *The Journal of Parasitology* 53:1055.
- Read, C. P., and M. Voge. 1954. The size attained by *Hymenolepis diminuta* in different host species. *The Journal of Parasitology* 40:88.
- Reynolds, L. A., K. J. Filbey, and R. M. Maizels. 2012. Immunity to the model intestinal helminth parasite *Heligmosomoides polygyrus*. *Seminars in immunopathology* 34:829–46.
- Roberts, L. 1961. The influence of population density on patterns and physiology of growth in *Hymenolepis diminuta* (Cestoda: Cyclophyllidea) in the definitive host. *Experimental Parasitology* 11:332–371.
- Rodriguez, R., and S. H. Kleven. 1980. Pathogenicity of two strains of *Mycoplasma gallisepticum* in broilers. *Avian Diseases* 24:800.
- Rogalski, M. A., and M. A. Duffy. 2020. Local adaptation of a parasite to solar radiation impacts disease transmission potential, spore yield, and host fecundity. *Evolution*.
- Rohr, J. R., D. J. Civitello, F. W. Halliday, P. J. Hudson, K. D. Lafferty, C. L. Wood, and E. A. Mordecai. 2020. Towards common ground in the biodiversity–disease debate. *Nature Ecology & Evolution* 4:24–33.
- Sakala, I. G., G. Chaudhri, A. A. Scalzo, P. Eldi, T. P. Newsome, R. M. Buller, and G. Karupiah. 2015. Evidence for persistence of ectromelia virus in inbred mice, recrudescence following immunosuppression and transmission to naïve mice. *PLoS Pathogens* 11:e1005342.
- Schindler, D. W. 1998. Whole-Ecosystem Experiments: Replication Versus Realism: The Need for Ecosystem-Scale Experiments. *Ecosystems* 1:323–334.
- Schmid-Hempel, P. 2011. *Evolutionary Parasitology: The Integrated Study of Infections, Immunology, Ecology, and Genetics* (1st ed.). Oxford University Press, Oxford.
- Schmid-Hempel, P., L. Wilfert, and R. Schmid-Hempel. 2019. Wildlife Diseases: linking theory to data. Pages 3–31 in K. Wilson, A. Fenton, and D. Thompkins, eds. *Pollinator diseases: the Bombus–Crithidia system*. Cambridge University Press, Cambridge, UK.
- Scott, M. E., and R. M. Anderson. 1984. The population dynamics of *Gyrodactylus bullatarudis* (Monogenea) within laboratory populations of the fish host *Poecilia reticulata*. *Parasitology* 89:159–194.

- Seppälä, O., A. Karvonen, and E. T. Valtonen. 2004. Parasite-induced change in host behaviour and susceptibility to predation in an eye fluke–fish interaction. *Animal Behaviour* 68:257–263.
- Seppälä, O., A. Karvonen, and E. T. Valtonen. 2005. Manipulation of fish host by eye flukes in relation to cataract formation and parasite infectivity. *Animal Behaviour* 70:889–894.
- Shykoff, J. A., and P. Schmid-Hempel. 1991. Parasites and the advantage of genetic variability within social insect colonies. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 243:55–58.
- Spence, P. J., W. Jarra, P. Lévy, A. J. Reid, L. Chappell, T. Brugat, M. Sanders, et al. 2013. Vector transmission regulates immune control of *Plasmodium* virulence. *Nature* 498:228–31.
- Stables, J., and L. Chappell. 1986a. Putative immune response of rainbow trout, *Salmo gairdneri*, to *Diplostomum spathaceum* infections. *Journal of Fish Biology* 29:115–122.
- Stables, J. N., and L. H. Chappell. 1986b. *Diplostomum spathaceum* (Rud. 1819): effects of physical factors on the infection of rainbow trout (*Salmo gairdneri*) by cercariae. *Parasitology* 93:71–79.
- Stephens, P. R., P. Pappalardo, S. Huang, J. E. Byers, M. J. Farrell, A. Gehman, R. R. Ghai, et al. 2017. Global Mammal Parasite Database version 2.0. *Ecology*, *Ecology* 98:1476–1476.
- Stevenson, M. M., J. J. Lyanga, and E. Skamene. 1982. Murine malaria: genetic control of resistance to *Plasmodium chabaudi*. *Infection and Immunity*.
- Tate, A. T., and A. L. Graham. 2017. Dissecting the contributions of time and microbe density to variation in immune gene expression. *Proceedings of the Royal Society B: Biological Sciences* 284:20170727–20170728.
- Taylor, L. H., M. J. Mackinnon, and A. F. Read. 1998. Virulence of mixed-clone and single-clone infections of the rodent malaria *Plasmodium chabaudi*. *Evolution* 58:583–591.
- Thrall, P. H., L. G. Barrett, P. N. Dodds, and J. J. Burdon. 2016. Epidemiological and Evolutionary Outcomes in Gene-for-Gene and Matching Allele Models. *Frontiers in Plant Science* 6:1084.
- Thrall, P. H., and J. J. Burdon. 2003. Evolution of virulence in a plant host-pathogen metapopulation. *Science* 299:1735–1737.
- Thrall, P. H., J. J. Burdon, and J. D. Bever. 2002. Local adaptation in the *Linum marginale*-*Melampsora Lini* host-pathogen interaction. *Evolution* 56:1340–1351.
- Travis, J. 2006. Is It What We Know or Who We Know? Choice of Organism and Robustness of Inference in Ecology and Evolutionary Biology. (American Society of Naturalists Presidential Address). *The American Naturalist* 167:303.

- Vidlak, D., and T. Kielian. 2016. Infectious dose dictates the host response during *Staphylococcus aureus* orthopedic-implant biofilm infection. *Infection and Immunity* 84:1957–1965.
- Voge, M. 1959a. Sensitivity of developing *Hymenolepis diminuta* larvae to high temperature stress. *The Journal of Parasitology* 45:175.
- . 1959b. Temperature stress and development of *Hymenolepis diminuta* in *Tribolium confusum* on different diets. *The Journal of Parasitology* 45:591.
- Voge, M., and D. Heyneman. 1958. Effect of high temperature on the larval development of *Hymenolepis nana* and *Hymenolepis diminuta* (Cestoda: Cyclophyllidea). *The Journal of Parasitology* 44:249.
- Wale, N., M. J. Jones, D. G. Sim, A. F. Read, and A. a King. 2019. The contribution of host cell-directed vs. parasite-directed immunity to the disease and dynamics of malaria infections. *Proceedings of the National Academy of Sciences of the United States of America* 2018:201908147.
- Wale, N., D. G. Sim, M. J. Jones, R. Salathe, T. Day, and A. F. Read. 2017. Resource limitation prevents the emergence of drug resistance by intensifying within-host competition. *Proceedings of the National Academy of Sciences of the United States of America* 114:13774–13779.
- Wargo, A. R., S. Huijben, J. C. de Roode, J. Shepherd, and A. F. Read. 2007. Competitive release and facilitation of drug-resistant parasites after therapeutic chemotherapy in a rodent malaria model. *Proceedings of the National Academy of Sciences of the United States of America* 104:19914–19919.
- Webster, J., S. Gowtage-Sequeira, M. Berdoy, and H. Hurd. 1999. Predation of beetles (*Tenebrio molitor*) infected with tapeworms (*Hymenolepis diminuta*): a note of caution for the Manipulation Hypothesis. *Parasitology* 120:313–318.
- Weimer, H. E., M. Voge, F. A. Quinn, and J. Redlich-Moshin. 1955. Serum glycoprotein concentrations in rats infected with the Cestode, *Hymenolepis diminuta*. *Proceedings of the Society for Experimental Biology and Medicine* 90:494–496.
- Weinstein, S. B., J. C. Buck, and H. S. Young. 2018. A landscape of disgust. *Science* 359:1213–1214.
- Whyte, S., L. Chappell, and C. Secombes. 1990. Protection of rainbow trout, *Oncorhynchus mykiss* (Richardson), against *Diplostomum spathaceum* (Digenea): the role of specific antibody and activated macrophages. *Journal of Fish Diseases* 13:281–291.
- Yan, G., L. Stevens, C. J. Goodnight, and J. J. Schall. 1998. Effects of a tapeworm parasite on the competition of *Tribolium*. *Ecology* 79:1093–1103.
- Yeung, F., Y.-H. Chen, J.-D. Lin, J. M. Leung, C. McCauley, J. C. Devlin, C. Hansen, et al. 2020. Altered Immunity of Laboratory Mice in the Natural Environment Is Associated with Fungal Colonization. *Cell Host & Microbe* 27:809-822.e6.

Zaiss, M. M., A. Rapin, L. Lebon, L. K. Dubey, I. Mosconi, K. Sarter, A. Piersigilli, et al. 2015. The intestinal microbiota contributes to the ability of helminths to modulate allergic inflammation. *Immunity* 43:998–1010.

Box 1: Key questions in EEID

EEID is a diverse field with roots in parasitology, immunology, epidemiology, and population biology in addition to ecology and evolutionary biology. It is no surprise, then, that the questions of interest are diverse. Here we outline some of the major themes of interest to EEID researchers.

Parasites as drivers of host population abundance, dynamics, and diversity (& vice versa)

‘(How) do parasites maintain genetic diversity in host populations?’ and ‘(how) do parasites control host population abundance?’ are two of the oldest questions in the field, the subject of papers that act as touchstones in the field and a continued subject of debate and research (Lively et al. 2014; Brandell et al. 2020). There has been longstanding interest in the reciprocal processes — how host population abundance and diversity impact the evolution of parasite traits, in particular virulence and transmissibility, and the coevolutionary dynamics that can thence ensue. Indeed, coevolutionary interactions have been of interest throughout EEID’s short history. Community level processes that could impact the probability of a parasite moving between hosts are a particularly active area of interest. In particular, the concept of the dilution effect, which states that host biodiversity at the community level can reduce the risk of disease in a particular host, is an ever-growing area of research (Keesing et al. 2010; Rohr et al. 2020).

The impact of the environment external to the host on disease

There is increasing interest in how ecological interactions, other than those between host and parasite, alter disease transmission and host-parasite (co)evolution (Lively et al. 2014). Perhaps

motivated by the pressing challenge of climate change and the prevalence of studies of vectorborne diseases in EEID (Brandell et al. 2020), the impact of temperature on disease dynamics has received considerable attention (Lafferty 2009; Mordecai et al. 2017). The effect of additional abiotic factors, including host diet and habitat structure, are also areas of active research (Hite et al. 2019). In addition to these abiotic factors, the role of predators in altering disease transmission, via their impact on host population density as well as through indirect effects, has been the subject of both theoretical and empirical research (Choo et al. 2003; Packer et al. 2003; Duffy et al. 2019). A notable recent trend is to understand how hosts select and use their habitats to avoid infection and how parasites, in turn, manipulate their hosts to promote their transmission (Hughes 2013; Weinstein et al. 2018).

The drivers of disease emergence

As infectious diseases emerge at an increasing rate (and at the time of writing, threaten our lives and livelihoods) (Jones et al. 2008), ecologists and evolutionary biologists have sought to understand the factors that drive their emergence. Emerging infectious diseases often spillover from animal reservoirs and, as a result, the ecological and evolutionary factors that enable parasites to 'jump' hosts and thence establish in a host population have received much attention (Lloyd-Smith et al. 2009; Babayan et al. 2018).

Within-host interactions

The impact of within-host interactions, whether that be between parasite strains/species or between parasites, hosts, and the microbiota, on disease dynamics is an active area of research that has perhaps the greatest overlap with the immunological and parasitological fields in which EEID is rooted. Studies of pathogen-pathogen interactions are conducted both to understand the eco-epidemiological dynamics of infectious diseases (e.g., how does a coinfection alter the spread of a focal pathogen? (Marchetto and Power 2018; Clay et al. 2019))

as well as to understand the evolution of parasite traits including virulence, transmissibility, and drug resistance (deRoode et al. 2005; Birger et al. 2015; Wale et al. 2017). Studies of the microbiota meanwhile focus on the microbiota as a unit (e.g., using antibiotics to ask 'how does the presence of microbiota alter host and pathogen fitness'), as well as its community ecology (e.g., 'how does the microbiota assemble?' 'How is it affected by pathogens & host genotype (and vice versa)?' And, 'are some species more important than others?') (Gonzalez et al. 2011; Koskella et al. 2017). Here, EEID overlaps with the wider field of microbial ecology. Within-host studies are often characterized by a focus on the on the dynamical nature of microbial populations; a major challenge in EEID is to understand how these dynamics translate to epidemic dynamics (Mideo et al. 2011; Clay et al. 2019).

Figure Legends

Figure 1: **The phenomena of interest to ecologists & evolutionary biologists of infectious diseases and the scales of biological organization at which they occur.** **A.** Epidemiological models define the central processes of importance to the ecology and evolution of infectious diseases. Individual hosts transition between epidemiologically distinct states (black): susceptible, exposed, infectious and recovered. Hosts transition between susceptible and infectious states via the process of transmission and exit via the process of recovery (or death, not shown). Both parasite (pink) & host (blue) traits are involved in the key phenomena (*italics*). On the part of the parasite, transmission is governed by infectivity (the parasite's capacity to successfully enter and establish within a host), replication (here defined, for simplicity, as the process by which parasites either replicate or mature within the host) and dispersal capacity, the ability to successfully exit the host. For the purposes of this review, when the entire cycle is completed and the parasite successfully enters a host, replicates, exits and makes it to a new host, we say that transmission is 'realized'. The host, of course, also determines parasite transmission: host susceptibility mediates the probability that a parasite successfully infects the host and the host can cause parasite mortality (our narrow definition of resistance). Infections cause disease (black box), to which both parasite and host may contribute, via virulence and immunopathology alike. Immunopathology and parasite virulence can be alleviated by host tolerance. Infectious hosts may exit the diseased state via the process of recovery, as mediated

by resistance, medications or death (not shown). The host and parasite traits involved in these phenomena may (co)evolve and are thus the focus of evolutionary studies of infectious diseases. Note that, in the case of multi-host parasites, each may trait may vary with the identity of the host. **B.** Infectious diseases impact, and are affected by, processes at multiple ecological scales. Parasite and host ecology and behavior can impact parasite (pink) and host (blue) traits and populations, and vice versa. In addition, other phenomena (black) at the within-host, population, community and ecosystem scales can impact parasite/host population dynamics and evolution, and vice versa. The challenge of model systems research is to investigate, and ideally quantify, the importance of these interactions for parasite ecology and evolution.

Figure 2: **The extent that model systems in EEID are used to investigate the key phenomena of interest to the field.** Top row: the shade of the square indicates the proportion of studies reviewed in which in the phenomenon is studied in our review. The number of study systems, of the 10 total, in which this phenomenon is the focus of study in at least once is indicated in the top right of the square. Bottom row: Each rectangle represents the trait or process indicated. Shading indicates the proportion of studies, which fall under the umbrella phenomenon, that focus on that trait/process e.g. 'morbidity' is bright orange, indicating that a large proportion of studies of disease focus on morbidity. Borders indicate whether the trait is a feature of the parasite (no border), host (solid) trait or neither (dashed). The numbers at the top of each rectangle indicate the number of study systems (of a total of 10) in which the focal trait is the focus of at least one study. Note that a single study can be allotted to more than one category.

Figure 3: **Model systems research tends to focus only on the small-scale ecological interactions that influence disease dynamics and there is limited vertical integration within model systems.** Each panel corresponds to one of the 10 systems reviewed. The scale of independent variables is indicated on the top of each panel, the scale of dependent variables on the left of each panel (see Table 3 for information on which experimental variables are allotted to each scale). White space indicates that we found no studies that focused on the interaction of processes at the corresponding scales in our review; shaded areas indicate that there was at least one. The intensity of shading indicates the proportion of studies in a certain system that fell into the corresponding category.

Table 1: **Experimental systems included in the literature review herein.**Superscripts indicate ¹first and ²second intermediate hosts, *definitive host.

Pathogen classifications follow those of Schmid-Hempel (2011).



Pathogen name(s)	Pathogen classification	Host classification	'Founding' EEID researcher(s) (decade of first publications)	Field of origin (% of 10 most cited experimental papers from EEID)	Life cycle & transmission mode	Can life cycle be completed in lab?	Recovery Occurs?
<i>Melampsora lini</i>	Microparasite, fungus	Plants	Henry Flor (1930s), Jeremy Burdon (1980s)	agriculture, genetics, (60%)	Direct; wind-borne	Yes	No, within a season (Ravensdalle et al. 2011)
<i>Infectious ectromelia virus</i>	Microparasite, virus	Rodent	Frank Fenner (1940s)	Medicine (smallpox model), EEID (0%)	Direct; physical contact, fomites	Yes	Yes (Chaudhri et al. 2004)
<i>Hymenolepis diminuta</i> or <i>Taenia taenia</i>	Macroparasite, cestode	Insects ¹ Rodent*	Marietta Voge (1950s), Clark Read, John Holmes	EEID (50%)	Indirect, obligate multi-host	No	No in rat; yes in mouse (Read 1967; Chappell and Pike 1976).
<i>Heligmosomoides polygyrus</i> or <i>Nematosporoides dubius</i>	Macroparasite, nematode	Mammals (rodents)	Clark Dobson (1960s), Anne Keymer	medicine, parasitology (10%)	Direct; ingestion of larvae.	Yes,	Yes (Reynolds et al. 2012)
<i>Diplostomum spathaceum</i>	Macroparasite, trematode	Molluscs ¹ Fish ²	John Stables Leslie Chappell (1990s), E. Tellervo Valtonen	fisheries, parasitology (100%)	Indirect, obligate multi-host	No. 1 st & 2 nd intermediate host studied	No (Whyte et al. 1990)
<i>Crithidia bombi</i>	Microparasite, protozoa	Arthropods (bumbees)	Paul Schmid-Hempel (1990s)	EEID (100%)	Direct; fecal-oral	Yes	Rarely (B. Sadd, <i>personal communication</i>)
<i>Pasteuria ramosa</i>	Microparasite, bacterium	Zooplankton	Dieter Ebert (1990s)	EEID (100%)	Direct; ingestion spores in water.	Yes	Rarely (Hall and Ebert 2012)
<i>Plasmodium chabaudi</i>	Macroparasite, protozoan	Rodents ¹ Arthropods*	Andrew Read (1990s)	Medicine (malaria model), parasitology (30%)	Indirect, vector-borne.	Yes, rarely.	Yes (Stevenson et al. 1982)
<i>Mycoplasma gallisepticum</i>	Microparasite, bacterium	Birds*	Andre Dhondt (1990s)	agriculture/ conservation (60%+)	Direct.	Yes	Yes (Kollias et al. 2004)
<i>Ribeiroia ondatrae</i>	Macroparasite, trematode	Molluscs ¹ Amphibians ² , Birds*	Pieter Johnson (1990s)	EEID/ conservation (100%)	Indirect, obligate multi-host	No. 1 st & 2 nd intermediate host studied	Varies with host (Johnson et al. 2004)

Table 2: **Definitions of categories to which studies were allotted.** Each study can be allotted to multiple categories.

phenomenon	organism	trait	definition
transmission	pathogen	infectivity	Capacity to successfully enter the host.
		replication	Capacity to grow in population size <i>or</i> grow/mature inside the host.
		dispersal capacity	Capacity to leave the host e.g. no. of transmission stages.
		entry to exit	Capacity to enter the host, replicate and be ready to transmit.
	realized transmission	Capacity to enter, replicate and exit a host and find a second.	
	host	susceptibility	Readiness with which host becomes infected
disease	pathogen	virulence	Contribution of parasites to host disease.
	host	tolerance	Capacity for host to maintain health when bearing a given burden of parasites.
		resistance	Capacity to kill parasites.

Table 3: **Definitions of scales to which variables were allotted.** Each variable within a study can be allotted to multiple categories.

Scale	trait	examples
within-host	Pertains to variation generated at, or processes that occur, within the host	<i>dynamics</i> of host or parasite cell types, presence or absence or number of co-infecting species or strains, tissue tropism, movement within host
individual	Pertains to a characteristic of an individual host or parasite or a host or parasite strain	genetics, behavior, sex, age, frequency, color, motility host only: immune status, maternal immune status parasite only: inoculum size,
ex-host	Abiotic aspect of the host environment that is proximate to the host individual	diet, light conditions, pH conditions, medication
population	Pertains to factors that vary at a population level. Can be generated due to temporal variation within a population or spatial variation among populations.	demography, age structure, social structure, genetic diversity.
community	Pertains to organisms other than the host or parasite or characteristics that define their interactions with them.	host diversity, predator presence or absence, antipredator behavior, host community composition, host range
ecosystem	Pertains to ecosystem scale processes and the measure that define them	chemical composition, biomass, nutrient cycling

A**B**


10 systems

transmission



10 systems

disease



7 systems

recovery




3 systems

other



% studies 25 50 75 100

scale of process varied



% studies 0 25 50 75